Inhibitors

MCE RedChemExpress

mPEG-amine (MW 20000)

Cat. No.: HY-140679 **CAS No.:** 80506-64-5


 $\label{eq:Molecular Formula:} \begin{array}{ll} \text{Molecular Formula:} & (C_{_2}H_{_4}O)_nC_{_3}H_{_9}NO \\ \\ \text{Target:} & \text{PROTAC Linkers} \end{array}$

Pathway: PROTAC

Storage: -20°C, protect from light, stored under nitrogen

* In solvent : -80°C, 6 months; -20°C, 1 month (protect from light, stored under

nitrogen)



BIOLOGICAL ACTIVITY

Description	m-PEG-NH2 (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] .
IC ₅₀ & Target	PEGs
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA