


## Azido-PEG2-alcohol

<b>Cat. No.:</b>	HY-140797		
<b>CAS No.:</b>	139115-90-5		
<b>Molecular Formula:</b>	C <sub>4</sub> H <sub>9</sub> N <sub>3</sub> O <sub>2</sub>		
<b>Molecular Weight:</b>	131.13		
<b>Target:</b>	PROTAC Linkers		
<b>Pathway:</b>	PROTAC		
<b>Storage:</b>	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

H<sub>2</sub>O : 100 mg/mL (762.60 mM; Need ultrasonic)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	7.6260 mL	38.1301 mL	76.2602 mL
5 mM	1.5252 mL	7.6260 mL	15.2520 mL
10 mM	0.7626 mL	3.8130 mL	7.6260 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

Azido-PEG2-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs<sup>[1]</sup>. Azido-PEG2-alcohol is a click chemistry reagent, it contains an Azide group and can undergo copper-catalyzed azide-alkyne cycloaddition reaction (CuAAC) with molecules containing Alkyne groups. Strain-promoted alkyne-azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.

#### IC<sub>50</sub> & Target

PEGs

#### In Vitro

PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins<sup>[1]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA