## Azido-PEG7-alcohol

Cat. No.: CAS No.: Molecular Formula: Molecular Weight: Target: Pathway: Storage:	HY-140798 1274892-60-2 C <sub>14</sub> H <sub>29</sub> N <sub>3</sub> O <sub>7</sub> 351.4 PROTAC Linkers PROTAC	<sup>*N</sup> *N* <sub>N</sub> ~~ <sup>0</sup> ~~ <sup>0</sup> ~~ <sup>0</sup> ~~ <sup>0</sup> ~~ <sup>0</sup>
---	---	--

BIOLOGICAL ACTIVITY		
Description	Azido-PEG7-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs <sup>[1]</sup> . Azido-PEG7-alcohol is a	
	click chemistry reagent, it contains an Azide group and can undergo copper-catalyzed azide-alkyne cycloaddition reaction (CuAAc) with molecules containing Alkyne groups. Strain-promoted alkyne-azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.	
IC <sub>50</sub> & Target	PEGs	
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

## REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

## Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Product Data Sheet

