N-(Azido-PEG3)-N-bis(PEG1-t-butyl ester)

| CAS No.: Molecular Formula: Molecular Weight: Target: Pathway: Storage: | HY-140873 2086689-00-9 C ₂₆ H ₄₈ N ₄ O ₁₀ 576.68 PROTAC Linkers PROTAC Please store the product under the recommended conditions in the Certificate of Analysis. | $\mathcal{M}_{\mathcal{M},\mathcal{M}} = \mathcal{M}_{\mathcal{M},\mathcal{M}} = \mathcal{M}_{\mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M} = \mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M},\mathcal{M},$ |
|--|---|---|
|--|---|---|

| BIOLOGICAL ACTIVITY | | |
|---------------------|---|--|
| BIOLOGICAL ACTIVITY | | |
| Description | N-(Azido-PEG3)-N-bis(PEG1-t-butyl ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] . N- (Azido-PEG3)-N-bis(PEG1-t-butyl ester) is a click chemistry reagent, it contains an Azide group and can undergo copper- catalyzed azide-alkyne cycloaddition reaction (CuAAc) with molecules containing Alkyne groups. Strain-promoted alkyne- azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups. | |
| IC₅₀ & Target | PEGs | |
| In Vitro | PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. | |

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA



Tel: 609-228-6898