


## Hexaethylene glycol

Cat. No.:	HY-141230		
CAS No.:	2615-15-8		
Molecular Formula:	C <sub>12</sub> H <sub>26</sub> O <sub>7</sub>		
Molecular Weight:	282.33		
Target:	PROTAC Linkers		
Pathway:	PROTAC		
Storage:	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### BIOLOGICAL ACTIVITY

Description	Hexaethylene glycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs <sup>[1]</sup> .
IC <sub>50</sub> & Target	PEGs
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA