


## SR1078

<b>Cat. No.:</b>	HY-14422		
<b>CAS No.:</b>	1246525-60-9		
<b>Molecular Formula:</b>	C <sub>17</sub> H <sub>10</sub> F <sub>9</sub> NO <sub>2</sub>		
<b>Molecular Weight:</b>	431.25		
<b>Target:</b>	ROR		
<b>Pathway:</b>	Metabolic Enzyme/Protease; Vitamin D Related/Nuclear Receptor		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	1 year
		-20°C	6 months



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 33.33 mg/mL (77.29 mM; Need ultrasonic)			
		Solvent Concentration	Mass	
			1 mg	5 mg
	<b>Preparing Stock Solutions</b>		10 mg	
	<b>1 mM</b>	2.3188 mL	11.5942 mL	23.1884 mL
	<b>5 mM</b>	0.4638 mL	2.3188 mL	4.6377 mL
	<b>10 mM</b>	0.2319 mL	1.1594 mL	2.3188 mL
Please refer to the solubility information to select the appropriate solvent.				
<b>In Vivo</b>	<ol style="list-style-type: none"> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 40% PEG300 &gt;&gt; 5% Tween-80 &gt;&gt; 45% saline Solubility: 2.5 mg/mL (5.80 mM); Suspended solution; Need ultrasonic</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (5.80 mM); Clear solution</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% corn oil Solubility: ≥ 2.5 mg/mL (5.80 mM); Clear solution</li> </ol>			

### BIOLOGICAL ACTIVITY

<b>Description</b>	SR1078 is a selective agonist of retinoic acid receptor-related orphan receptor α/γ (RORα/RORγ). SR1078 directly binds to the ligand binding domain of RORα and RORγ and increases the transcriptional activity of these receptors, leading to stimulation of RORα/γ target gene transcription <sup>[1][2]</sup> .
<b>IC<sub>50</sub> &amp; Target</b>	RORα/γ <sup>[1]</sup>
<b>In Vitro</b>	SR1078 is a synthetic RORα/RORγ ligand. SR1078 modulates the conformation of RORγ in a biochemical assay and activates

ROR $\alpha$  and ROR $\gamma$  driven transcription. Furthermore, SR1078 stimulates expression of endogenous ROR target genes in HepG2 cells that express both ROR $\alpha$  and ROR $\gamma$ . In a cell-based chimeric receptor Gal4 DNA-binding domain-NR ligand binding domain cotransfection assay, SR1078 significantly inhibits the constitutive transactivation activity of ROR $\alpha$  and ROR $\gamma$ , but has no effect on the activity of FXR, LXR $\alpha$  and LXR $\beta$ . In a ROR $\alpha$  cotransfection assay, treatment of cells with SR1078 (10  $\mu$ M) results in a significant increase in transcription. Similarly, in the ROR $\gamma$  cotransfection assay, SR1078 treatment results in a stimulation of ROR $\gamma$ -dependent transcription activity<sup>[1]</sup>.

?SR1078 (2-10  $\mu$ M; 24 hours) shows a dose-dependent increase in expression of A2BP1, CYP19A1, NLGN1, and IPTR1 in SH-SY5Y cells. EC50's are in the range of 3-5  $\mu$ M<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

The pharmacokinetic properties of SR1078 are examined in mice and noted significant exposure. Plasma concentrations reach 3.6  $\mu$ M 1h after a 10 mg/kg i.p. injection of SR1078 and sustained levels of above 800 nM even 8h after the single injection. These levels are sufficient to perform a proof-of-principle experiment to determine if SR1078 treatment would stimulate ROR target gene expression in an animal model. Mice are treated with SR1078 (10 mg/kg; i.p.) and 2h after the injection the livers are harvested and mRNA purified for assessment of G6Pase and FGF21 gene expression. The expression of both FGF21 and G6Pase is significantly stimulated by SR1078 treatment vs. vehicle control<sup>[1]</sup>.

?SR1078 (10 mg/kg; i.p.) shows a significant 25% reduction in repetitive grooming behavior in the BTBR mice<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

#### Kinase Assay <sup>[1]</sup>

The ALPHA screen assays are performed. Assays are performed in triplicate in white opaque 384-well plates under green light conditions (<100 lux) at room temperature. The final assay volume is 20  $\mu$ L. All dilutions are made in assay buffer (100 mM NaCl, 25 mM HEPES, 0.1% (w/v) BSA, pH 7.4). The final DMSO concentration is 0.25% (v/v). A mix of 12  $\mu$ L of GST-ROR $\gamma$ -LBD (10 nM), beads (12.5  $\mu$ g/mL of each donor and acceptor), and 4  $\mu$ L of increasing concentrations (210 nM-50  $\mu$ M) of compound SR1078 are added to the wells, the plates are sealed and incubated for 1h. After this preincubation step, 4  $\mu$ L of Biotin-TRAP220-2 peptide (50 nM) is added, the plates are sealed and further incubated for 2h. The plates are read on PerkinElmer Envision 2104 and data analyzed using GraphPad Prism software<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Cell Assay <sup>[1]</sup>

HEK293 cells are maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% (v/v) fetal bovine serum at 37°C under 5% CO<sub>2</sub>. HepG2 cells are maintained and routinely propagated in minimum essential medium supplemented with 10% (v/v) fetal bovine serum at 37°C under 5% CO<sub>2</sub>. In experiments where lipids and sterols are depleted, cells are maintained on charcoal treated serum (10% (v/v) fetal bovine serum) and treated with 7.5  $\mu$ M lovastatin and 100  $\mu$ M mevalonic acid. 24 h prior to transfection, HepG2 or HEK293 cells are plated in 96-well plates at a density of 15 $\times$ 10<sup>3</sup> cells/well. Transfections are performed using Lipofectamine<sup>TM</sup> 2000. 16 h post-transfection, the cells are treated with vehicle or SR1078. 24 h post-treatment, the luciferase activity is measured using the Dual-Glo<sup>TM</sup> luciferase assay system. The experiments are repeated at least three times<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Animal Administration <sup>[1]</sup>

Mice<sup>[1]</sup>

Plasma levels of SR1078 are evaluated in C57BL6 mice (n=3 per time point) administered by i.p. injection. After 1, 2, 4, and 8h blood is taken. In the 2h time point liver is taken for target gene analysis. Plasma is generated using standard centrifugation techniques, and the plasma and tissues are frozen at -80°C. Plasma and tissues are mixed with acetonitrile (1:5 (v/v) or 1:5 (w/v), respectively), sonicated with a probe tip sonicator, and analyzed for drug levels by liquid chromatography/tandem mass spectrometry.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Acta Pharm Sin B. 2018, 53(1): 62-67.
- Hepatology. 2021 Sep 12.
- Cell Death Differ. 2023 Apr 7.
- Cell Death Dis. 2021 Sep 28;12(10):886.
- NPJ Parkinsons Dis. 2022 Jul 8;8(1):90.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

---

- [1]. Wang Y, et al. Identification of SR1078, a synthetic agonist for the orphan nuclear receptors ROR $\alpha$  and ROR $\gamma$ . ACS Chem Biol. 2010 Nov 19;5(11):1029-34.
- [2]. Wang Y, Billon C, Walker JK, Burris TP. Therapeutic Effect of a Synthetic ROR $\alpha$ / $\gamma$  Agonist in an Animal Model of Autism. ACS Chem Neurosci. 2016;7(2):143-148.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA