


Phospholipid-PEG-Biotin (MW 3400)

Cat. No.:	HY-147207B
Target:	Liposome
Pathway:	Metabolic Enzyme/Protease
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description

Phospholipid-PEG-Biotin (MW 3400) is a phospholipid PEG derivative that has a biotin and a phospholipid bridged by a linear PEG linker. Phospholipid-PEG-Biotin (MW 3400) can interact with avidinylated antibodies. Phospholipid-PEG-Biotin (MW 3400) can be used to modify liposome and cells surface, and pancreatic islets for cell transplantation^{[1][2][3]}.

REFERENCES

- [1]. Chen Y, et al. A rapid and efficient technique for liposomal and nonliposomal drug pharmacokinetics studies using magnetic nanoprobes and its application to leakage kinetics of liposomes. *J Chromatogr A*. 2018 Dec 14;1580:2-11.
- [2]. Hoenius M, et al. Biotinylated Stealth magnetoliposomes. *Chem Phys Lipids*. 2002 Dec;120(1-2):75-85.
- [3]. Teramura Y, et al. Influence of molecular weight of PEG chain on interaction between streptavidin and biotin-PEG-conjugated phospholipids studied with QCM-D. *Acta Biomater*. 2016 Jan;30:135-143.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA