


ZK53

Cat. No.:	HY-149677		
Molecular Formula:	C ₁₉ H ₂₀ BrF ₂ N ₃ O		
Molecular Weight:	424.28		
Target:	Mitochondrial Metabolism		
Pathway:	Metabolic Enzyme/Protease		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 200 mg/mL (471.39 mM; ultrasonic and warming and heat to 80°C)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	2.3569 mL	11.7847 mL	23.5693 mL
5 mM	0.4714 mL	2.3569 mL	4.7139 mL
10 mM	0.2357 mL	1.1785 mL	2.3569 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: 5 mg/mL (11.78 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: 5 mg/mL (11.78 mM); Clear solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

ZK53 is a selective activator of mitochondrial caseinolytic protease P (HsClpP) (EC₅₀: 1.37?μM for α-casein hydrolysis by HsClpP). ZK53 is inactive toward bacterial ClpP proteins. ZK53 induces apoptosis in H1703, H520 and SK-MES-1 cells. ZK53 induces dysregulation of mitochondrial functions in lung squamous cell carcinoma (LUSC) cells. ZK53 inhibits tumor growth in H1703 xenograft mouse model^[1].

REFERENCES

- [1]. Zhou LL, et al. Selective activator of human ClpP triggers cell cycle arrest to inhibit lung squamous cell carcinoma. Nat Commun. 2023 Nov 3;14(1):7069.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA