Screening Libraries

Inhibitors

•

•

Proteins

Steroid sulfatase/17 β -HSD1-IN-3

Cat. No.:	HY-151201	
CAS No.:	2990549-76-1	
Molecular Formula:	C ₁₉ H ₁₇ ClN ₂ O ₅ S	
Molecular Weight:	420.87	
Target:	Steroid Sulfatase	
Pathway:	Metabolic Enzyme/Protease	
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	

BIOLOGICAL ACTIVITY		
Description	Steroid sulfatase/17β-HSD1-IN-3 (compound 19) is a dual inhibitor of steroid sulfatase (STS) and 17β-hydroxysteroid dehydrogenase type 1 (17β HSD1). Steroid sulfatase/17β-HSD1-IN-3 irreversibly inhibits hSTS activity with anIC ₅₀ value of 27 nM. Steroid sulfatase/17β-HSD1-IN-3 can be used in the study of endometriosis and other estrogen-dependent diseases ^[1] .	
IC ₅₀ & Target	IC50: 27 nM (hSTS) ^[1] .	
In Vitro	Steroid sulfatase/17β-HSD1-IN-3 (compound 19) (0-1100 nM, 24 h) inhibits human steroid sulfatase (hSTS) with an IC ₅₀ value of 27 nM and acts on irreversible inhibition of hSTS with an IC ₅₀ value of 29 nM in T47D human breast cancer cells ^[1] . Steroid sulfatase/17β-HSD1-IN-3 (compound 19) (20 μM, 48 h) inhibits HEK-293 cells growth by 30% at a concentration of 20 μM with low cytotoxicity ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

REFERENCES

[1]. Abdelrahman Mohamed, et al. Dual Targeting of Steroid Sulfatase and 17β-Hydroxysteroid Dehydrogenase Type 1 by a Novel Drug-Prodrug Approach: A Potential Therapeutic Option for the Treatment of Endometriosis. J Med Chem. 2022 Aug 22.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

