


(R)-(+)-Bay-K-8644

Cat. No.:	HY-15125		
CAS No.:	98791-67-4		
Molecular Formula:	C ₁₆ H ₁₅ F ₃ N ₂ O ₄		
Molecular Weight:	356.3		
Target:	Calcium Channel		
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 300 mg/mL (841.99 mM)
 H₂O : < 0.1 mg/mL (insoluble)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		2.8066 mL	14.0331 mL	28.0662 mL
	5 mM		0.5613 mL	2.8066 mL	5.6132 mL
	10 mM		0.2807 mL	1.4033 mL	2.8066 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: ≥ 2.5 mg/mL (7.02 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
 Solubility: 2.5 mg/mL (7.02 mM); Suspended solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

(R)-(+)-Bay-K-8644 is a calcium channel inhibitor. (R)-(+)-Bay-K-8644 inhibits Ba²⁺ currents (I_{Ba}) (IC₅₀=975 nM).

IC₅₀ & Target

IC₅₀: 975 nM (I_{Ba})^[1]

In Vitro

(R)-(+)-Bay-K-8644, a conventional racemic mixture of Bay K 8644, is widely used as an L-type Ca²⁺ channel agonist. Each optical isomer possesses opposite effects on I_{Ba}, (R)-(+)-Bay-K-8644 as an antagonist and S(-)-Bay K 8644 as an agonist. (R)-(+)-Bay-K-8644 inhibits Ba²⁺ currents (I_{Ba}) (IC₅₀=975 nM). When (R)-(+)-Bay-K-8644 (0.5 μM) is applied, I_{Ba} is suppressed to 71±10% of control. In the presence of (R)-(+)-Bay-K-8644^[1], (R)-(+)-Bay-K-8644 is a calcium channel inhibitor^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Zhu HL, et al. Antagonistic actions of S(-)-Bay K 8644 on cyclic nucleotide-induced inhibition of voltage-dependent Ba(2+) currents in guinea pig gastric antrum. *Naunyn Schmiedebergs Arch Pharmacol.* 2008 Dec;378(6):609-15.
- [2]. Sidaway P, et al. L-type Ca²⁺ channel sparklets revealed by TIRF microscopy in mouse urinary bladder smooth muscle. *PLoS One.* 2014 Apr 3;9(4):e93803.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA