


## SARS-CoV-2-IN-26

<b>Cat. No.:</b>	HY-151269
<b>Molecular Formula:</b>	C <sub>52</sub> H <sub>52</sub> O <sub>8</sub> P <sub>2</sub>
<b>Molecular Weight:</b>	866.91
<b>Target:</b>	SARS-CoV
<b>Pathway:</b>	Anti-infection
<b>Storage:</b>	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

<b>Description</b>	SARS-CoV-2-IN-23 is a two-armed diphosphate ester and medium length molecular tweezers. SARS-CoV-2-IN-23 exhibits antiviral activity with IC <sub>50</sub> s of 8.2 μM and 2.6 μM against SARS-CoV-2 activity and the spike pseudoparticle transduction, respectively. SARS-CoV-2-IN-23 induces liposomal membrane disruption with an EC <sub>50</sub> value of 4.4 μM <sup>[1]</sup> .								
<b>IC<sub>50</sub> &amp; Target</b>	IC <sub>50</sub> : 4.4 μM (viral liposome, SARS-CoV-2) <sup>[1]</sup>								
<b>In Vitro</b>	<p>SARS-CoV-2-IN-23 (CP002) inhibits SARS-CoV-2 (IC<sub>50</sub>=2.6 μM) with a low cytotoxicity (Caco2 cells, CC<sub>50</sub>=97 μM)<sup>[1]</sup>. SARS-CoV-2-IN-23 (0-15 μM; 2 h) inactivate SARS-CoV-2, shows inhibition against infection with an IC<sub>50</sub> value of 8.2 μM<sup>[1]</sup>. SARS-CoV-2-IN-23 suppresses various enveloped viruses activity with IC<sub>50</sub>s of 38.4 μM (influenza A virus, IAV), 2.8 μM (measles virus, MeV), 1.1 μM (herpes simplex viruses, HSV-1), and human immunodeficiency virus type 1 (HIV-1), respectively<sup>[1]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Cell Viability Assay<sup>[1]</sup></p> <table border="1"> <tr> <td>Cell Line:</td> <td>Caco2 cells exposed with SARS-CoV-2 (2 h, 37 °C)</td> </tr> <tr> <td>Concentration:</td> <td>0, 0.23, 0.93, 3.75, 15 μM</td> </tr> <tr> <td>Incubation Time:</td> <td>2 hours; determined infection rates on day 2</td> </tr> <tr> <td>Result:</td> <td>Inhibited SARS-CoV-2 infection activity to Caco2 cells.</td> </tr> </table>	Cell Line:	Caco2 cells exposed with SARS-CoV-2 (2 h, 37 °C)	Concentration:	0, 0.23, 0.93, 3.75, 15 μM	Incubation Time:	2 hours; determined infection rates on day 2	Result:	Inhibited SARS-CoV-2 infection activity to Caco2 cells.
Cell Line:	Caco2 cells exposed with SARS-CoV-2 (2 h, 37 °C)								
Concentration:	0, 0.23, 0.93, 3.75, 15 μM								
Incubation Time:	2 hours; determined infection rates on day 2								
Result:	Inhibited SARS-CoV-2 infection activity to Caco2 cells.								

### REFERENCES

[1]. Tatjana Weil, et al. Advanced Molecular Tweezers with Lipid Anchors against SARS-CoV-2 and Other Respiratory Viruses. JACS Au 2022, XXXX, XXX, XXX-XXX.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA