(Me)Tz-butanoic acid

Cat. No.: CAS No.: Molecular Formula: Molecular Weight: Target: Pathway: Storage:	HY-151730 1923268-81-8 C_7H_10N_4O_2 182.18 ADC Linker Antibody-drug Conjugate/ADC Related Please store the product under the recommended conditions in the Certificate of Analysis.	N.N.O N.N.OH
---	--	-----------------

	ITV																						
OLOGICAL ACTIV	ΙΤΥ)	-																				
Description	(Me)Tz	<u>'</u> -	bu	butano	butanoic acid	butanoic acid is a cli	butanoic acid is a click ch	butanoic acid is a click chemist	butanoic acid is a click chemistry reag	butanoic acid is a click chemistry reagent of	butanoic acid is a click chemistry reagent contain	butanoic acid is a click chemistry reagent containing a	butanoic acid is a click chemistry reagent containing an azic	butanoic acid is a click chemistry reagent containing an azide. (M	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-l	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-butan	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-butanoic ac	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-butanoic acid is	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-butanoic acid is tetra	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-butanoic acid is tetrazine	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-butanoic acid is tetrazine linker	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-butanoic acid is tetrazine linker for	butanoic acid is a click chemistry reagent containing an azide. (Me)Tz-butanoic acid is tetrazine linker for
	conjug	52	atic	ation via	ation via Diels-	ation via Diels-Alder-	ation via Diels-Alder-react	ation via Diels-Alder-reaction, i.	ation via Diels-Alder-reaction, i.e. cop	ation via Diels-Alder-reaction, i.e. copper-fr	ation via Diels-Alder-reaction, i.e. copper-free Clio	ation via Diels-Alder-reaction, i.e. copper-free Click rea	ation via Diels-Alder-reaction, i.e. copper-free Click reaction	ation via Diels-Alder-reaction, i.e. copper-free Click reaction with	ation via Diels-Alder-reaction, i.e. copper-free Click reaction with dieno	ation via Diels-Alder-reaction, i.e. copper-free Click reaction with dienophiles	ation via Diels-Alder-reaction, i.e. copper-free Click reaction with dienophiles $^{[1]}$. (M	ation via Diels-Alder-reaction, i.e. copper-free Click reaction with dienophiles $^{[1]}$. (Me)Tz	tion via Diels-Alder-reaction, i.e. copper-free Click reaction with dienophiles ^[1] . (Me)Tz-buta	ation via Diels-Alder-reaction, i.e. copper-free Click reaction with dienophiles $^{[1]}$. (Me)Tz-butanoic	ation via Diels-Alder-reaction, i.e. copper-free Click reaction with dienophiles $^{[1]}$. (Me)Tz-butanoic acid i	tion via Diels-Alder-reaction, i.e. copper-free Click reaction with dienophiles ^[1] . (Me)Tz-butanoic acid is a clic	ation via Diels-Alder-reaction, i.e. copper-free Click reaction with dienophiles ^[1] . (Me)Tz-butanoic acid is a click
	chemi	s	try	try reag	try reagent, it	try reagent, it contai	try reagent, it contains a ⊺	try reagent, it contains a Tetraz	try reagent, it contains a Tetrazine gro	try reagent, it contains a Tetrazine group tl	try reagent, it contains a Tetrazine group that car	try reagent, it contains a Tetrazine group that can unde	try reagent, it contains a Tetrazine group that can undergo a	try reagent, it contains a Tetrazine group that can undergo an inv	try reagent, it contains a Tetrazine group that can undergo an inverse e	try reagent, it contains a Tetrazine group that can undergo an inverse electro	try reagent, it contains a Tetrazine group that can undergo an inverse electron der	try reagent, it contains a Tetrazine group that can undergo an inverse electron demand	try reagent, it contains a Tetrazine group that can undergo an inverse electron demand Diel	try reagent, it contains a Tetrazine group that can undergo an inverse electron demand Diels-Ald	try reagent, it contains a Tetrazine group that can undergo an inverse electron demand Diels-Alder rea	try reagent, it contains a Tetrazine group that can undergo an inverse electron demand Diels-Alder reaction (try reagent, it contains a Tetrazine group that can undergo an inverse electron demand Diels-Alder reaction (iED
	with m	10	ole	olecules	olecules conta	olecules containing	olecules containing TCO g	olecules containing TCO groups	plecules containing TCO groups.	plecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.	blecules containing TCO groups.

REFERENCES

[1]. Sadowsky Jack Avid, et al. METHODS AND COMPOSITIONS FOR TIME DELAYED BIO-ORTHOGONAL RELEASE OF CYTOTOXIC AGENTS. WO2022197877. 2022

Caution: Product has not been fully validated for medical applications. For research use only.

Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Product Data Sheet



Tel: 609-228-6898