


## PF-04691502

<b>Cat. No.:</b>	HY-15177		
<b>CAS No.:</b>	1013101-36-4		
<b>Molecular Formula:</b>	C <sub>22</sub> H <sub>27</sub> N <sub>5</sub> O <sub>4</sub>		
<b>Molecular Weight:</b>	425.48		
<b>Target:</b>	PI3K; mTOR; Autophagy		
<b>Pathway:</b>	PI3K/Akt/mTOR; Autophagy		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 50 mg/mL (117.51 mM; Need ultrasonic)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	<b>Preparing Stock Solutions</b>	1 mM	2.3503 mL	11.7514 mL	23.5029 mL
		5 mM	0.4701 mL	2.3503 mL	4.7006 mL
10 mM		0.2350 mL	1.1751 mL	2.3503 mL	
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	<ol style="list-style-type: none"> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 40% PEG300 &gt;&gt; 5% Tween-80 &gt;&gt; 45% saline Solubility: ≥ 2.5 mg/mL (5.88 mM); Clear solution</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (5.88 mM); Clear solution</li> <li>Add each solvent one by one: 10% DMSO &gt;&gt; 90% corn oil Solubility: ≥ 2.5 mg/mL (5.88 mM); Clear solution</li> </ol>				

### BIOLOGICAL ACTIVITY

<b>Description</b>	PF-04691502 is a potent and selective inhibitor of PI3K and mTOR. PF-04691502 binds to human PI3Kα, β, δ, γ and mTOR with K <sub>i</sub> s of 1.8, 2.1, 1.6, 1.9 and 16 nM, respectively.			
<b>IC<sub>50</sub> &amp; Target</b>	PI3Kδ 1.6 nM (K <sub>i</sub> )	PI3Kα 1.8 nM (K <sub>i</sub> )	PI3Kγ 1.9 nM (K <sub>i</sub> )	PI3Kβ 2.1 nM (K <sub>i</sub> )
	mTOR 16 nM (K <sub>i</sub> )			

<b>In Vitro</b>	<p>PF-04691502 inhibits recombinant mouse PI3K<math>\alpha</math> in an ATP-competitive inhibitor. PF-04691502 potently inhibits AKT phosphorylation on S473 and T308 in all the 3 cancer cell lines with IC<sub>50</sub> values of 3.8 to 20 nM and 7.5 to 47 nM, respectively. Using a 96-well plate-based P-S6RP(S235/236) ELISA assay, PF-04691502 potently inhibits mTORC1 activity with an IC<sub>50</sub> of 32 nM. PF-04691502 inhibits cell proliferation of BT20, SKOV3, and U87MG with IC<sub>50</sub> values of 313, 188, and 179 nM, respectively. In PIK3CA-mutant and PTEN-deleted cancer cell lines, PF-04691502 reduces phosphorylation of AKT T308 and AKT S473 (IC<sub>50</sub> of 7.5-47 nM and 3.8-20 nM, respectively) and inhibits cell proliferation (IC<sub>50</sub> of 179-313 nM). PF-04691502 inhibits mTORC1 activity in cells as measured by PI3K-independent nutrient stimulated assay, with an IC<sub>50</sub> of 32 nM and inhibits the activation of PI3K and mTOR downstream effectors including AKT, FKHRL1, PRAS40, p70S6K, 4EBP1, and S6RP<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
<b>In Vivo</b>	<p>Nude mice bearing U87MG tumors are administered orally once a day with PF-04691502 at 0.5, 1, 5, and 10 mg/kg (maximum tolerated dose, MTD). Treatment with 10 mg/kg results in a significant reduction of P-AKT(S473) levels at 1 hour postdosing, and persistent inhibition is observed for 8 hours. P-AKT(S473) recovers to above baseline 24 hours after 10 mg/kg treatment. For P-S6RP(S235/236), a similar inhibition time course is observed, but after 24 hours of treatment, P-S6RP levels remain lower than vehicle tumors. Modulation of the AKT downstream effector, P-PRAS40(T246), and mTOR downstream effector, P-4EBP1(T37/46), is observed. The PF-04691502-treated tumors are also evaluated by immunohistochemistry for levels of P-AKT(S473), total AKT, P-S6RP, and total S6RP. Phosphorylation of AKT and S6RP are significantly reduced at 4 hours after a single dose of PF-04691502 at 10 mg/kg. Dose-dependent tumor growth inhibition (TGI) is obtained in the U87MG xenograft model and approximately 73% TGI is observed at the MTD dose of 10 mg/kg<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

## PROTOCOL

<b>Kinase Assay</b> <sup>[1]</sup>	<p>The biochemical protein kinase assays for class I PI3K and mTOR are assessed. The fluorescence polarization assay for ATP competitive inhibition is done as follows: mPI3K<math>\alpha</math> dilution solution (90 nM) is prepared in fresh assay buffer (50 mM Hepes pH 7.4, 150 mM NaCl, 5 mM DTT, 0.05% CHAPS) and kept on ice. The enzyme reaction contained 0.5 nM mouse PI3K<math>\alpha</math> (p110 <math>\alpha</math>/p85<math>\alpha</math> complex purified from insect cells), 30 <math>\mu</math>M PIP2, PF-04691502 (0, 1, 4, and 8 nM), 5 mM MgCl<sub>2</sub>, and 2-fold serial dilutions of ATP (0-800 <math>\mu</math>M). Final DMSO is 2.5%. The reaction is initiated by the addition of ATP and terminated after 30 minutes with 10 mM EDTA. In a detection plate, 15 <math>\mu</math>L of detector/probe mixture containing 480 nM GST-Grp1PH domain and 12 nM TAMRA tagged fluorescent PIP3 in assay buffer is mixed with 15 <math>\mu</math>L of kinase reaction mixture. The plate is shaken for 3 minutes, and incubated for 35 to 40 minutes before reading on an LJL Analyst HT<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
<b>Cell Assay</b> <sup>[1]</sup>	<p>BT20, U87MG, and SKOV3 cells are plated at 3,000 cell/well in 96-well culture plates in growth medium with 10% FBS. Cells are incubated overnight and treated with DMSO (0.1% final) or serial diluted compound for 3 days. Resazurin is added to 0.1 mg/mL. Plates are incubated at 37°C in 5% CO<sub>2</sub> for 3 hours. Fluorescence signals are read as emission at 590 nm after excitation at 530 nm. IC<sub>50</sub> values are calculated by plotting fluorescence intensity to drug concentration in nonlinear curves. U87MG and SKOV3 cells are plated in 96-well plates overnight and caspase-3/caspase-7 activity is assessed with the Caspase-Glo 3/7 Assay Kit<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
<b>Animal Administration</b> <sup>[1]</sup>	<p>Mice<sup>[1]</sup></p> <p>Female nu/nu mice (6-8 weeks old) are used. Tumor cells for implantation are harvested and resuspended in serum-free medium mixed with matrigel (1:1). SKOV3, U87MG, or NSCLC cells (2.5-4<math>\times</math>10<sup>6</sup>) are implanted subcutaneously into the hind flank region. Treatment started when average tumor size is 100 to 200 mm<sup>3</sup>. PF-04691502 is formulated in 0.5% methylcellulose in water suspension and given orally once a day. Animal body weights and tumor volumes are measured every 2 to 3 days. Tumor volume is determined with Vernier calipers and calculated. Percentage of tumor growth inhibition (TGI) is calculated. Data are presented as mean<math>\pm</math>SE. Comparisons between treatment groups and vehicle group are done using 1-way ANOVA by Dunnett's tests. Student's t test is used to determine the P value for the comparison of 2 groups.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

---

## CUSTOMER VALIDATION

- Sci Transl Med. 2018 Jul 18;10(450):eaaq1093.
- Cell Death Differ. 2021 Jul;28(7):2221-2237.
- Theranostics. 2020 Jan 1;10(4):1531-1543.
- Molecules. 2020 Apr 23;25(8):1980.
- bioRxiv. 2020 Jun.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

---

## REFERENCES

[1]. Yuan J, et al. PF-04691502, a potent and selective oral inhibitor of PI3K and mTOR kinases with antitumor activity. Mol Cancer Ther. 2011 Nov;10(11):2189-99.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA