


## GSK 650394

<b>Cat. No.:</b>	HY-15192		
<b>CAS No.:</b>	890842-28-1		
<b>Molecular Formula:</b>	C <sub>25</sub> H <sub>22</sub> N <sub>2</sub> O <sub>2</sub>		
<b>Molecular Weight:</b>	382.45		
<b>Target:</b>	SGK; Influenza Virus		
<b>Pathway:</b>	Metabolic Enzyme/Protease; Anti-infection		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 40.7 mg/mL (106.42 mM)  
 \* "≥" means soluble, but saturation unknown.

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	2.6147 mL	13.0736 mL	26.1472 mL
5 mM	0.5229 mL	2.6147 mL	5.2294 mL
10 mM	0.2615 mL	1.3074 mL	2.6147 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
 Solubility: 2.5 mg/mL (6.54 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
 Solubility: ≥ 2.5 mg/mL (6.54 mM); Suspended solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
 Solubility: ≥ 2.5 mg/mL (6.54 mM); Clear solution

### BIOLOGICAL ACTIVITY

#### Description

GSK 650394 is a novel SGK inhibitor with IC<sub>50</sub> of 62 nM and 103 nM for SGK1 and SGK2 in the SPA assay respectively. GSK 650394 also inhibits influenza virus replication.

#### IC<sub>50</sub> & Target

SGK1

#### In Vitro

GSK650394 is relatively non-toxic, with LC<sub>50</sub> values of 41 μM in M1 cells (68 times its activity IC<sub>50</sub>) and a LC<sub>50</sub> greater than 100

$\mu\text{M}$  in HeLa cells. GSK650394 inhibits SGK1-mediated epithelial transport with an  $\text{IC}_{50}$  of  $0.6 \mu\text{M}$  in the SCC assay. GSK650394 inhibits the growth of LNCaP cells with  $\text{IC}_{50}$  of approximately  $1 \mu\text{M}$ <sup>[1]</sup>. GSK650394A inhibits the insulin-induced phosphorylation of PKB-Ser<sup>473</sup> at  $3 \mu\text{M}$ , and essentially abolishes this response at  $10 \mu\text{M}$ . GSK650394A ( $1\text{-}10 \mu\text{M}$ ) does not alter the phosphorylation of PRAS40-Ser246 in hormone-deprived cells or prevent the insulin-induced phosphorylation of this residue<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

GSK650394 ( $1, 10, \text{ and } 30 \mu\text{M}, 10 \mu\text{L}/\text{rat}$ , intrathecally) dose-dependently prevents CFA-induced pain behavior and the associated SGK1 phosphorylation, GluR1 trafficking, and protein-protein interactions at 1 day after CFA administration<sup>[3]</sup>. GSK650394 at concentrations of  $10, 30, \text{ and } 100 \text{ nM}$  ( $10 \mu\text{L}$ ), but not vehicle solution (SNL 3D+Veh and SNL 7D+Veh, respectively), dose-dependently increases the withdrawal latency of the ipsilateral hindpaw at 1-3 and 1-5 h after injection at days 3 and 7 postsurgery (SNL 3D+GSK and SNL 7D+GSK, respectively). GSK650394 (from day 0 to 6 postsurgery;  $100 \text{ nM}, 10 \mu\text{L}$ , i.t.) administration alleviates SNL-induced allodynia at days 3, 5, and 7 postsurgery in SNL animals<sup>[4]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

#### Cell Assay <sup>[1]</sup>

The toxicity of GSK650394 to M-1 and HeLa cells is assessed using the Cell Proliferation Kit (XTT) following manufacturer's instructions. Briefly,  $10,000$  HeLa or M-1 cells/well are plated into 96-well plates in  $100 \mu\text{L}$  of the appropriate maintenance media. After 48 h, media is removed and replaced with  $100 \mu\text{L}$  of EMEM with Earle's salts containing  $2 \text{ mM}$  L-glutamine and  $1\%$  antibiotic-antimycotic overnight. M-1 cells are also supplemented with  $1 \mu\text{g}/\text{mL}$  insulin,  $6.25 \mu\text{g}/\text{mL}$  sodium selenite, and  $6.25 \mu\text{g}/\text{mL}$  transferrin. After 24 h, the media is removed and replaced with  $100 \mu\text{L}$  media alone or media containing increasing concentrations of GSK650394. For HeLa cells,  $50 \mu\text{L}$  of activated XTT solution is added after 4 h. For M-1 cells,  $50 \mu\text{L}$  of activated XTT solution is added after 24 h. Following a 2 h incubation, absorbance is measured at  $490 \text{ nm}$  using a SpectraMAX PLUS spectrophotometer and the data analyzed to obtain  $\text{IC}_{50}$  values using GraphPad Prism 3 software. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Animal Administration <sup>[4]</sup>

Briefly, the rats are anesthetized under isoflurane anesthesia (induction  $5\%$ , maintenance  $2\%$  in oxygen). An incision is made, and the left L5 spinal nerves are carefully isolated and tightly ligated with 6-0 silk sutures  $2\text{-}5 \text{ mm}$  distal to the dorsal root ganglia. GSK650394 ( $10, 30, \text{ and } 100 \text{ nM}, 10 \mu\text{L}$ ) is administered by bolus injection at 3 or 7 d or by daily injection for 7 d (day 0-6) postspinal nerve ligation. A vehicle solution of a volume identical to that of the tested agents is dispensed to serve as a control.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Sci Transl Med. 2018 Jul 18;10(450):eaaq1093.
- Int J Biol Sci. 2023; 19(1): 204-224.
- Oncogene. 2021 Sep;40(35):5367-5378.
- Biomed Pharmacother. 2024 Mar 21;174:116447.
- J Transl Med. 2023 Aug 14;21(1):544.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Sherk AB, et al. Development of a small-molecule serum- and glucocorticoid-regulated kinase-1 antagonist and its evaluation as a prostate cancer therapeutic. Cancer

---

Res. 2008 Sep 15;68(18):7475-83.

[2]. Mansley MK, et al. Effects of nominally selective inhibitors of the kinases PI3K, SGK1 and PKB on the insulin-dependent control of epithelial Na<sup>+</sup> absorption. *Br J Pharmacol.* 2010 Oct;161(3):571-88.

[3]. Peng HY, et al. Spinal SGK1/GRASP-1/Rab4 is involved in complete Freund's adjuvant-induced inflammatory pain via regulating dorsal horn GluR1-containing AMPA receptor trafficking in rats. *Pain.* 2012 Dec;153(12):2380-92.

[4]. Peng HY, et al. Spinal serum-inducible and glucocorticoid-inducible kinase 1 mediates neuropathic pain via kalirin and downstream PSD-95-dependent NR2B phosphorylation in rats. *J Neurosci.* 2013 Mar 20;33(12):5227-40.

[5]. Alamares-Sapuay JG, et al. Serum- and glucocorticoid-regulated kinase 1 is required for nuclear export of the ribonucleoprotein of influenza A virus. *J Virol.* 2013 May;87(10):6020-6.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA