


## N7-Methyl-2'-O-(2-methoxyethyl) guanosine

Cat. No.:	HY-152568
Molecular Formula:	C <sub>14</sub> H <sub>21</sub> N <sub>5</sub> O <sub>6</sub>
Molecular Weight:	355.35
Target:	Nucleoside Antimetabolite/Analog
Pathway:	Cell Cycle/DNA Damage
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

#### Description

N7-Methyl-2'-O-(2-methoxyethyl) guanosine is a purine nucleoside analog. Purine nucleoside analogs have broad antitumor activity targeting indolent lymphoid malignancies. Anticancer mechanisms in this process rely on inhibition of DNA synthesis, induction of apoptosis, etc<sup>[1]</sup>.

### REFERENCES

- [1]. Virág L, Szabó C. Purines inhibit poly(ADP-ribose) polymerase activation and modulate oxidant-induced cell death. *FASEB J.* 2001 Jan;15(1):99-107.
- [2]. Saugstad OD. Hypoxanthine as an indicator of hypoxia: its role in health and disease through free radical production. *Pediatr Res.* 1988 Feb;23(2):143-50.
- [3]. Robak T, Robak P. Purine nucleoside analogs in the treatment of rarer chronic lymphoid leukemias. *Curr Pharm Des.* 2012;18(23):3373-88.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA