MCE MedChemExpress

Product Data Sheet

9-(β-D-Xylofuranosyl)guanine


Cat. No.:HY-152673CAS No.:27462-39-1Molecular Formula: $C_{10}H_{13}N_5O_5$ Molecular Weight:283.24

Target: Nucleoside Antimetabolite/Analog

Pathway: Cell Cycle/DNA Damage

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.



BIOLOGICAL ACTIVITY

Description

9-(β-D-Xylofuranosyl)guanine is a guanosine analog. Some guanosine analogs have immunostimulatory activity. In some animal models, they also induce type I interferons, producing antiviral effects. Studies have shown that the functional activity of guanosine analogs is dependent on the activation of Toll-like receptor 7 (TLR7)^[1].

REFERENCES

[1]. Robak T, Robak P. Purine nucleoside analogs in the treatment of rarer chronic lymphoid leukemias. Curr Pharm Des. 2012;18(23):3373-88.

[2]. Lee J, et al. Molecular basis for the immunostimulatory activity of guanine nucleoside analogs: activation of Toll-like receptor 7. Proc Natl Acad Sci U S A. 2003 May 27;100(11):6646-51.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA