Caspase-3/7 activator 3

Cat. No.:	HY-152758	\wedge
Molecular Formula:	C ₂₄ H ₂₇ NO ₅	
Molecular Weight:	409.47	
Target:	Apoptosis; Caspase	
Pathway:	Apoptosis	$\langle \rangle O H_{N}$
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	

BIOLOGICAL ACTIVITY				
Description	Caspase-3/7 activator 3 is an effective activator of Caspase-3/7. Caspase-3/7 activator 3 has tumor selectivity, anti- proliferative activity and high Apoptosis-inducing ability ^[1] .			
IC ₅₀ & Target	Caspase-3	Caspase-7		
In Vitro	Caspase-3/7 activator 3 (Compound 7g) inhibits Wi-38, Caco-2 and HepG-2 cells with IC ₅₀ values of 17.077 μM, 0.064 μM, 0.065 μM, respectively ^[1] . Caspase-3/7 activator 3 (0.06 μM; 72 h) significantly inhibits the expression of apoptosis inhibitor Bcl2 in Caco-2 cells (down-regulates Bcl2 by 0.087 folds). Caspase-3/7 activator 3 insignificantly effects the morphology of Caco-2 and HepG-2 cells ^[1] . Caspase-3/7 activator 3 (0.064 μM; 72 h) has the potential to induces apoptosis of Caco-2 cells (the total apoptotic cell populations is 58.730 %) ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. Apoptosis Analysis ^[1]			
	Cell Line:	Caco-2 cells.		
	Concentration:	0.064 μΜ.		
	Incubation Time:	72 h.		
	Result:	Showed to promote cell apoptosis.		

REFERENCES

Page 1 of 2

[1]. Ayoup MS, et al. Structure optimization of new tumor-selective Passerini α-acyloxy carboxamides as Caspase-3/7 activators. Sci Rep. 2022 Dec 27;12(1):22390.

Product Data Sheet



Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA