## RedChemExpress

## Product Data Sheet

## 5-(2-Hydroxyethyl)-2',3'-di-O-toluoyl-2'-deoxyuridine

Cat. No.:HY-154160Molecular Formula:C2,7 H2,8 N2 O8Molecular Weight:508.52Target:Nucleoside Antimetabolite/AnalogPathway:Cell Cycle/DNA DamageStorage:Please store the product under the recommended conditions in the Certificate of Analysis.			
Molecular Weight:       508.52         Target:       Nucleoside Antimetabolite/Analog         Pathway:       Cell Cycle/DNA Damage         Storage:       Please store the product under the recommended conditions in the Certificate of	Cat. No.:	HY-154160	
Target:       Nucleoside Antimetabolite/Analog         Pathway:       Cell Cycle/DNA Damage         Storage:       Please store the product under the recommended conditions in the Certificate of	Molecular Formula:	$C_{27}H_{28}N_{2}O_{8}$	0
Pathway:       Cell Cycle/DNA Damage         Storage:       Please store the product under the recommended conditions in the Certificate of	Molecular Weight:	508.52	HN HN
Storage: Please store the product under the recommended conditions in the Certificate of	Target:	Nucleoside Antimetabolite/Analog	
	Pathway:	Cell Cycle/DNA Damage	
	Storage:		

BIOLOGICAL ACTIVITY			
Description	5-(2-Hydroxyethyl)-2',3'-di-O-toluoyl-2'-deoxyuridine is a purine nucleoside analog. Purine nucleoside analogs have broad antitumor activity targeting indolent lymphoid malignancies. Anticancer mechanisms in this process rely on inhibition of DNA synthesis, induction of apoptosis, etc <sup>[1]</sup> .		

## REFERENCES

[1]. Robak T, Robak P. Purine nucleoside analogs in the treatment of rarer chronic lymphoid leukemias. Curr Pharm Des. 2012;18(23):3373-88.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

.OH