


BML-190

Cat. No.:	HY-15420		
CAS No.:	2854-32-2		
Molecular Formula:	C ₂₃ H ₂₃ ClN ₂ O ₄		
Molecular Weight:	426.89		
Target:	Cannabinoid Receptor		
Pathway:	GPCR/G Protein; Neuronal Signaling		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (117.13 mM; Need ultrasonic)					
		Solvent Concentration	Mass	1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM		2.3425 mL	11.7126 mL	23.4252 mL
		5 mM		0.4685 mL	2.3425 mL	4.6850 mL
10 mM			0.2343 mL	1.1713 mL	2.3425 mL	
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (5.86 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	<p>BML-190(IMMA) is a potent and selective CB2 receptor ligand (K_i values are 435 nM and > 2 μM for CB2 and CB1 respectively). IC₅₀ Value: 435 nM(K_i CB2)Target:CB2 receptor in vitro: BML-190 increases the accumulation of cAMP, via forskolin-stimulated mechanism in HEK-293 cells. Alternate studies suggest that BML-190 reduces the toxicity of culture supernatants to SH-SY5Y human neuroblastoma cells. Various research suggests that BML-190 is an essential tool in studying the proliferation of neuroblastoma. BML-190 diminishes LPS-induced NO and IL-6 production in a concentration-dependent manner. BML-190 also inhibits LPS-induced PGE2 production and COX-2 induction. in vivo:</p>
--------------------	--

REFERENCES

[1]. Zhang, Qiang; Ma, Peng; Cole, Richard B.; Wang, Guangdi In vitro metabolism of indomethacin morpholinylamide (BML-190), an inverse agonist for the peripheral cannabinoid receptor (CB2) in rat liver microsomes. European Journal of Pharmaceutical Sciences (

[2]. Klegeris, Andis; Bissonnette, Christopher J.; McGeer, Patrick L. Reduction of human monocytic cell neurotoxicity and cytokine secretion by ligands of the cannabinoid-type CB2 receptor. *British Journal of Pharmacology* (2003), 139(4), 775-786.

[3]. New DC, Wong YH. BML-190 and AM251 act as inverse agonists at the human cannabinoid CB2 receptor: signalling via cAMP and inositol phosphates. *FEBS Lett.* 2003 Feb 11;536(1-3):157-60.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA