


Salubrial

Cat. No.:	HY-15486
CAS No.:	405060-95-9
Molecular Formula:	C ₂₁ H ₁₇ Cl ₃ N ₄ OS
Molecular Weight:	479.8
Target:	Phosphatase; Autophagy; Apoptosis; HSV
Pathway:	Metabolic Enzyme/Protease; Autophagy; Apoptosis; Anti-infection
Storage:	4°C, protect from light * In solvent : -80°C, 2 years; -20°C, 1 year (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (104.21 mM; Need ultrasonic)				
		Mass			
		Solvent			
		Concentration			
Preparing Stock Solutions	1 mM	2.0842 mL	10.4210 mL	20.8420 mL	
	5 mM	0.4168 mL	2.0842 mL	4.1684 mL	
	10 mM	0.2084 mL	1.0421 mL	2.0842 mL	
	Please refer to the solubility information to select the appropriate solvent.				
In Vivo	1. Add each solvent one by one: 45% PEG300 >> 5% Tween-80 >> 50% saline Solubility: 10 mg/mL (20.84 mM); Suspended solution; Need ultrasonic				

BIOLOGICAL ACTIVITY

Description	Salubrial is a cell-permeable and selective inhibitor of eIF2α dephosphorylation ^[1] . Salubrial acts as a dual-specificity phosphatase 2 (Dusp2) inhibitor and suppresses inflammation in anti-collagen antibody-induced arthritis ^[2] . Salubrial has antiviral activity against HSV-1 and inhibits dephosphorylation of eIF2α mediated by the HSV-1 protein ICP34.5 ^[3] .	
IC₅₀ & Target	Dusp2	HSV-1
In Vitro	Salubrial, a recently identified PP1 inhibitor capable to protect against endoplasmic reticulum (ER) stress in various model systems, strongly synergized with proteasome inhibitors to augment apoptotic death of different leukemic cell lines. Salubrial preferentially seems to target the PP1/GADD34 complex, Salubrial is of interest to examine whether the effect of Salubrial could also be recapitulated by another inhibitor of this phosphatase. For this purpose cantharidin, was selected, which is less toxic than okadaic acid, but which also blocks PP1 (IC ₅₀ =1.7 μM) activities ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
In Vivo	Salubrial is a synthetic chemical that inhibits de-phosphorylation of eukaryotic translation initiation factor 2 alpha (eIF2α).	

Salubrinal significantly suppresses inflammation of the paws of CAIA mice. For instance, the clinical scores are 1.94 ± 1.7 (placebo) and 0.31 ± 0.6 (Salubrinal) on day 6; and 4.63 ± 3.4 (placebo) and 1.09 ± 1.6 (Salubrinal) on day 12. Consistent with the clinical scores, the thickening of the paws is also reduced in the Salubrinal-treated group. Furthermore, Salubrinal reduces the histological scores from 1.47 ± 1.10 (N=16; placebo) to 0.59 ± 0.64 (N=16; Salubrinal) ($p=0.01$)^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay ^[1]

Phosphatase activities are determined on immunoprecipitates of the phosphatases. Briefly, 2×10^6 K562 cells are treated for 18 hr with Salubrinal (20 μ M), PSI (10 nM), the combination of both drugs or okadaic acid (100 nM). After washing with PBS, cells are lysed for 15 min on ice either in PP1LB (for determination of PP1 γ -activity; 20 mM Tris-HCl, pH 7.5, 1% Triton X-100, 10% glycerol, 132 mM NaCl, Roche complete protease inhibitor) or in RIPA (for PP2A), supplemented with Roche complete protease inhibitor). Cell lysates containing 500 μ g (PP1 γ) or 300 μ g (PP2A) protein are immunoprecipitated overnight at 4°C with 2-3 μ g of the appropriate antibodies and then incubated with Protein A-Sepharose. Immunoprecipitates are washed three times in lysis buffer, followed by resuspension in phosphatase assay buffer (PP2A: 20 mM Tris-HCl, pH 7.5, 0.1 mM CaCl₂; PP1 γ : 50 mM Tris HCl pH 7.0, 0.2 mM MnCl₂, 0.1 mM CaCl₂, 125 μ g/mL BSA, 0.05% Tween 20), supplemented with 100 μ M 6,8-difluoro-4-methyl-umbelliferyl phosphate (DiFMUP). Precipitates are allowed to react with substrate for 1 hr at 37°C on an Eppendorf Thermoshaker, centrifuged and DiFMU fluorescence is measured on a BioTek Lambda Fluoro 320 microplate reader (360 nm_{ex}/460 nm_{em}). Phosphatase activities are given as percent change relative to the control (DMSO treated cells) ^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Assay ^[1]

Cellular viability is assessed by the WST-1 colorimetric assay. Assays are performed on 96 well plates with 2×10^4 K562 cells/well in triplicate with Salubrinal concentrations ranging from 5-75 μ M (total volume of 200 μ L, 18 hrs). Untreated cells served as negative control sample^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[2]

Mice^[2]

Using Balb/c female mice (~nine weeks old), CAIA is induced by intravenous injection of a 2 mg cocktail of ArthritoMAB antibodies on day 0 followed by intraperitoneal injection of 100 μ g LPS on day 3. Mice are randomly divided into a placebo group and a Salubrinal-treated group. Salubrinal (2.0 mg/kg) is intravenously administered daily from day 0, while a solvent (49.5% PEG 400 and 0.5% Tween 80 in PBS) is administered to the placebo group.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Nature. 2023 Sep;621(7977):188-195.
- Acta Biomater. 2020 Jun;109:229-243.
- Cell Prolif. 2021 Sep 28:e13133.
- EMBO Rep. 2022 Apr 11:e53932.
- Biomed Pharmacother. 2022 Dec 14;158:114133.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Drexler HC. Synergistic Apoptosis Induction in Leukemic Cells by the Phosphatase Inhibitor Salubrinal and Proteasome Inhibitors. PLoS One. 2009;4(1):e4161.

[2]. Hamamura K, et al. Salubrinal acts as a Dusp2 inhibitor and suppresses inflammation in anti-collagen antibody-induced arthritis. *Cell Signal*. 2015 Apr;27(4):828-35.

[3]. Bryant KF, et al. ICP34.5-dependent and -independent activities of salubrinal in herpes simplex virus-1 infected cells. *Virology*. 2008 Sep 30;379(2):197-204.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA