18:1 PEG3000 PE

MedChemExpress

Molecular Formula: $(C_2H_4O)_nC_{43}H_{80}NO_{10}P.NH_3$ Target: Liposome Pathway: Metabolic Enzyme/Protease Storage: Please store the product under the recommended conditions in the Certificate of Analysis.	ни ⁹ го~1°, орбон состо во нь
--	--

BIOLOGICAL ACTIVITY	
Description	18:1 PEG3000 PE is a PEG lipid functional end group used in the synthesis of liposomes (LPs) for the design of conjugated polymer nanoparticles. Through biotin modification and carboxyl terminus, lipid nanoparticles (LNPs) further coupling with other biomolecules can be achieved. Functionalized nanoparticles can be used for targeted labeling of specific cellular proteins. With streptavidin as a linker, biotinylated PEG lipid-conjugated polymer nanoparticles are able to bind to biotinylated antibodies on cell surface receptors, yielding the utility of fluorescence-based imaging and sensing.

REFERENCES

[1]. And resen TL, et al. Compositional inhomogeneity of drug delivery liposomes quantified at the single liposome level. Acta Biomater. 2020 Dec;118:207-214.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA