Product Data Sheet

N-Nonanoylglycine-d₂

 $\begin{array}{lll} \textbf{Cat. No.:} & & \text{HY-158084S} \\ \textbf{CAS No.:} & & 3008541\text{-}78\text{-}1 \\ \textbf{Molecular Formula:} & & C_{11}\text{H}_{19}\text{D}_{2}\text{NO}_{3} \\ \end{array}$

Molecular Weight: 217.3

Target: Isotope-Labeled Compounds

Pathway: Others

Storage: Powder -20°C 3 years

4°C 2 years

In solvent -80°C 6 months

-20°C 1 month

N O D D O

BIOLOGICAL ACTIVITY

 $\label{eq:Description} \textbf{N-Nonanoylglycine-} d_2 \text{ is the deuterated labeled N-Nonanoylglycine} [1].$

In Vitro Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as

tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to

affect the pharmacokinetic and metabolic profiles of $drugs^{[1]}$.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Screening Libraries

Inhibitors

1 000