


Irinotecan

Cat. No.:	HY-16562
CAS No.:	97682-44-5
Molecular Formula:	C ₃₃ H ₃₈ N ₄ O ₆
Molecular Weight:	586.68
Target:	Topoisomerase; Autophagy
Pathway:	Cell Cycle/DNA Damage; Autophagy
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (42.61 mM; Need ultrasonic)						
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg	
				1 mM	1.7045 mL	8.5225 mL	17.0451 mL
				5 mM	0.3409 mL	1.7045 mL	3.4090 mL
				10 mM	0.1705 mL	0.8523 mL	1.7045 mL
Please refer to the solubility information to select the appropriate solvent.							
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (3.55 mM); Clear solution						
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (3.55 mM); Clear solution						
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (3.55 mM); Clear solution						

BIOLOGICAL ACTIVITY

Description	Irinotecan ((+)-Irinotecan) is a topoisomerase I inhibitor, preventing religation of the DNA strand by binding to topoisomerase I-DNA complex ^[1] .
IC ₅₀ & Target	Topoisomerase I
In Vitro	Irinotecan is a topoisomerase I inhibitor. Irinotecan inhibits the growth of LoVo and HT-29 cells, with IC ₅₀ s of 15.8 ± 5.1 and 5.17 ± 1.4 μM, respectively, and induces similar amounts of cleavable complexes in both in LoVo and HT-29 cells ^[2] . Irinotecan suppresses the proliferation of human umbilical vein endothelial cells (HUVEC), with an IC ₅₀ of 1.3 μM ^[3] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Irinotecan (CPT-11, 5 mg/kg) significantly inhibits the growth of tumors by intratumoral injection daily for 5 days, on two consecutive weeks in rats, and such effects also occur via continuous intraperitoneal infusion by osmotic minipump into mice. However, Irinotecan (10 mg/kg) shows no effect on the growth of tumor by i.p.^[1] Irinotecan (CPT-11, 100-300 mg/kg, i.p.) apparently suppresses tumor growth of HT-29 xenografts in athymic female mice by day 21. The two groups of Irinotecan (125 mg/kg) plus TSP-1 (10 mg/kg per day) or Irinotecan (150 mg/kg) in combination TSP-1 (20 mg/kg per day) are nearly equally effective and inhibit tumor growth 84% and 89%, respectively, and both are more effective than Irinotecan alone at doses of 250 and 300 mg/kg^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[2]

Exponentially growing cells are seeded in 20 cm² dishes with an optimal cell number for each cell line (20,000 for LoVo cells, 100,000 for HT-29 cells). They are treated 2 days later with increasing concentrations of irinotecan or SN-38 for one cell doubling time (24 h for LoVo cells, 40 h for HT-29 cells). After washing with 0.15 M NaCl, the cells are further grown for two doubling times in normal medium, detached from the support with trypsin-EDTA and counted in a hemocytometer. The IC₅₀ values are then estimated as the drug concentrations responsible for 50% growth inhibition as compared with cells incubated without drug^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[1]

Irinotecan has been administered by intratumoral injection at 0.1 cc volume of the appropriate solution, for a doses of 5 mg/kg daily for 5 days, on two consecutive weeks, followed by a 7-days rest period, referred to as one cycle of therapy. Rats receive three cycles over a period of 8 weeks. Control animals receive 0.1 cc of sterile 0.9% sodium chloride solution by intratumoral injection in the same rule of administration as that of animals of group II^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Genome Med. 2016 Oct 31;8(1):116.
- Theranostics. 2019 May 31;9(13):3732-3753.
- Cell Death Dis. 2019 Nov 25;10(12):887.
- PLoS Pathog. 2020 Mar 24;16(3):e1008429.
- Pharmacol Res. 2020 Oct 4;105232.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Morales C, et al. Antitumoral effect of irinotecan (CPT-11) on an experimental model of malignant neuroectodermal tumor. *J Neurooncol.* 2002 Feb;56(3):219-26.
- [2]. Pavillard V, et al. Determinants of the cytotoxicity of irinotecan in two human colorectal tumor cell lines. *Cancer Chemother Pharmacol.* 2002 Apr;49(4):329-35. Epub 2002 Jan 30.
- [3]. Allegrini G, et al. Thrombospondin-1 plus irinotecan: a novel antiangiogenic-chemotherapeutic combination that inhibits the growth of advanced human colon tumor xenografts in mice. *Cancer Chemother Pharmacol.* 2004 Mar;53(3):261-6. Epub 2003 Dec 5.
- [4]. Dela Cruz FS, et al. A case study of an integrative genomic and experimental therapeutic approach for rare tumors: identification of vulnerabilities in a pediatric poorly differentiated carcinoma. *Genome Med.* 2016 Oct 31;8(1):116.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA