


## Apabetalone

Cat. No.:	HY-16652	
CAS No.:	1044870-39-4	
Molecular Formula:	C <sub>20</sub> H <sub>22</sub> N <sub>2</sub> O <sub>5</sub>	
Molecular Weight:	370.4	
Target:	Epigenetic Reader Domain; HIV	
Pathway:	Epigenetics; Anti-infection	
Storage:	Powder	-20°C 3 years
		4°C 2 years
	In solvent	-80°C 1 year
		-20°C 6 months



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 33 mg/mL (89.09 mM)  
 \* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	2.6998 mL	13.4989 mL	26.9978 mL
	5 mM	0.5400 mL	2.6998 mL	5.3996 mL
	10 mM	0.2700 mL	1.3499 mL	2.6998 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 50% PEG300 >> 50% saline  
Solubility: 15.15 mg/mL (40.90 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 2.5 mg/mL (6.75 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
Solubility: ≥ 2.5 mg/mL (6.75 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
Solubility: ≥ 2.5 mg/mL (6.75 mM); Clear solution
- Add each solvent one by one: 5% DMSO >> 40% PEG300 >> 5% Tween-80 >> 50% saline  
Solubility: ≥ 2.5 mg/mL (6.75 mM); Clear solution
- Add each solvent one by one: 5% DMSO >> 95% (20% SBE-β-CD in saline)  
Solubility: ≥ 2.5 mg/mL (6.75 mM); Clear solution
- Add each solvent one by one: 1% DMSO >> 99% saline  
Solubility: ≥ 0.5 mg/mL (1.35 mM); Clear solution

## BIOLOGICAL ACTIVITY

<b>Description</b>	Apabetalone (RVX-208) is an inhibitor of BET transcriptional regulators with selectivity for the second bromodomain. The IC <sub>50</sub> s are 87 μM and 0.51 μM for BD1 and BD2, respectively <sup>[1]</sup> .
<b>IC<sub>50</sub> &amp; Target</b>	IC50: 510±41 nM (BD2), 87±10 μM (BD1) <sup>[1]</sup>
<b>In Vitro</b>	Apabetalone (RVX-208) competes with binding of an acetylated histone peptide to tandem BD1 BD2 protein constructs of the four BET proteins, with IC <sub>50</sub> s between 0.5 and 1.8 μM. Apabetalone increases the production of ApoA-I in hepatocytes in vitro, which results in increased high density lipoprotein cholesterol (HDL-C). Apabetalone selectively binds to bromodomains of the BET (Bromodomain and Extra Terminal) family, competing for a site bound by the endogenous ligand, acetylated lysine, and that this accounts for its pharmacological activity. Apabetalone increases Apolipoprotein A-I (ApoA-I) production through an epigenetic mechanism and suggests that BET inhibition may be a promising new approach to the treatment of atherosclerosis. Apabetalone increases ApoA-I expression in liver cells <sup>[2]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
<b>In Vivo</b>	In the atherosclerosis prophylactic treatment study design, mice are fed a Western diet concurrent with the treatment with 150 mg/kg/dose b.i.d. for 12 weeks. Mice are sacrificed at 12 weeks after treatment. There is a progressive increase in body weight in both the vehicle treated as well as the Apabetalone (RVX-208) treated groups. However, there is only an increase of 4 g (from 24 g to 28 g) body weight after 12 weeks on Western diet in the Apabetalone treated group whereas this increase is found to be 9 g (25 g-34 g) in the vehicle treated group. The significant decrease in body weight gain in Apabetalone treated mice is not due to decreased feed consumption, suggesting a positive attribute of the molecule. Plasma lipid measurements are done at 6 weeks and 12 weeks of treatment with either the vehicle or Apabetalone. Compared to the vehicle control animals, Apabetalone treated mice show significant increase (~200%) in the levels of HDL-C at 6 weeks of treatment, which is sustained until end of the study (12 weeks) <sup>[3]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

<b>Cell Assay</b> <sup>[2]</sup>	Huh7 cells are plated at 23,000/well in a 96 well plate in DMEM+10% FBS before allowing to grow overnight. Cells are treated with compounds for 48 h in 0.1% DMSO with or without 5 μM Actinomycin D. U937 cells are differentiated for 3 days in 60 ng/mL PMA, 32,000 cells/well in 96-well format. Cells are then treated with compound in 0.1% DMSO in RPMI media+10% FBS, and after 1 h, lipopolysaccharide is added to the cells at 1 μg/mL for 3 hours <sup>[2]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
<b>Animal Administration</b> <sup>[3]</sup>	Mice <sup>[3]</sup> Seven to eight week old male ApoE <sup>-/-</sup> mice are used. Based on the body weight and lipid values, mice are divided into 2 groups (n=12): group 1, vehicle; and group 2, test agent, Apabetalone. Mice are then switched to Western diet (0.15% cholesterol and 42% calories from fat) and concurrently treated orally by gavage with either vehicle or the test agent, Apabetalone (150 mg/kg/dose b.i.d) for 12 weeks. After 6 week of treatment, an interim blood draw is done to monitor serum lipid levels. After 12 weeks of treatment mice are sacrificed to measure blood lipid parameters, aortic lesion, and liver and aortic RNA. Eight mice are used for enface (aortic plaque) analysis, 4 mice for tissue collection for mRNA and all 12 mice used for aortic sinus lesion area measurement. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Cell. 2021 Apr 15;184(8):2167-2182.e22.
- Elife. 2020 Dec 7;9:e61405.

- Antioxid Redox Signal. 2021 Dec 16.
- J Mol Cell Cardiol. 2018 Dec 7;127:83-96.
- Patent. US20180263995A1.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

---

## REFERENCES

- [1]. Picaud S, et al. RVX-208, an inhibitor of BET transcriptional regulators with selectivity for the second bromodomain. Proc Natl Acad Sci U S A. 2013 Dec 3;110(49):19754-9.
- [2]. McLure KG, et al. RVX-208, an inducer of ApoA-I in humans, is a BET bromodomain antagonist. PLoS One. 2013 Dec 31;8(12):e83190.
- [3]. Jahagirdar R, et al. A novel BET bromodomain inhibitor, RVX-208, shows reduction of atherosclerosis in hyperlipidemic ApoE deficient mice. Atherosclerosis. 2014 Sep;236(1):91-100.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA