


Ezetimibe

Cat. No.:	HY-17376		
CAS No.:	163222-33-1		
Molecular Formula:	C ₂₄ H ₂₁ F ₂ NO ₃		
Molecular Weight:	409.43		
Target:	Keap1-Nrf2; Autophagy		
Pathway:	NF-κB; Autophagy		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 200 mg/mL (488.48 mM; Need ultrasonic)
 H₂O : < 0.1 mg/mL (ultrasonic;warming;heat to 60°C) (insoluble)

Preparing Stock Solutions	Solvent / Mass		1 mg	5 mg	10 mg
	Concentration				
	1 mM		2.4424 mL	12.2121 mL	24.4242 mL
	5 mM		0.4885 mL	2.4424 mL	4.8848 mL
	10 mM		0.2442 mL	1.2212 mL	2.4424 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: 5 mg/mL (12.21 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% corn oil
 Solubility: ≥ 2.5 mg/mL (6.11 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Ezetimibe (SCH 58235) is a potent cholesterol absorption inhibitor. Ezetimibe is a Niemann-Pick C1-like1 (NPC1L1) inhibitor, and is a potent Nrf2 activator.

IC₅₀ & Target

NPC1L1, Nrf2^[1]

In Vitro

Ezetimibe (Eze) acts as a potent Nrf2 activator without causing cytotoxicity. Ezetimibe enhances transactivation of Nrf2, as revealed by a luciferase reporter assay. Ezetimibe also upregulates Nrf2 target genes, including GSTA1, heme oxygenase-1 (HO-1) and Nqo-1 in Hepa1c1c7 and MEF cells. Ezetimibe upregulates Nrf2 target genes in Nrf2^{+/+} MEF cells, whereas this induction is totally blocked in Nrf2^{-/-} MEF cells. Taken together, Ezetimibe acts as a novel Nrf2 inducer in a ROS-independent

manner^[1]. Human huh7 hepatocytes are pretreated with Ezetimibe (10 μ M, 1 h) and incubated with palmitic acid (PA, 0.5 mM, 24 h) to induce hepatic steatosis. Ezetimibe treatment significantly attenuates PA-increased triglycerides (TG) levels, which is consistent with our animal study. PA treatment resulted in an approximately 20% decrease in mRNA expression of ATG5, ATG6, and ATG7, which had been increased by Ezetimibe treatment. In addition, Ezetimibe treatment significantly increased the PA-induced reduction in LC3 protein abundance^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Administration of Ezetimibe (Eze) reduces the liver weights of mice fed the methionine- and choline-deficient (MCD) diet. This is consistent with the beneficial effects of Ezetimibe on hepatic steatosis. Liver histology shows pronounced multiple macrovesicular fat droplets in mice on the MCD diet, but Ezetimibe treatment markedly decreases the number and size of those droplets. Furthermore, hepatic fibrosis in mice fed the MCD diet is significantly attenuated by Ezetimibe^[1]. Blood and liver lipid levels including TG, free fatty acids (FFA), and total cholesterol (TC) are significantly decreased in Ezetimibe-treated OLETF rats. Moreover, OLETF rats show higher serum levels of glucose, insulin, HOMA-IR, TG, FFA, and TC than LETF animals, which are significantly reduced by Ezetimibe. In addition, histological analysis indicated that OLETF control rats showed larger lipid droplets in hepatocytes than age-matched LETO controls, which are attenuated by administration of Ezetimibe^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay ^[1]

GST-p62 is prepared from Escherichia coli and 0.5 μ g of the purified GST-p62 protein is used for in vitro AMPK phosphorylation assay. Phosphorylation of p62 protein by AMPK is determined by non-radioisotope method using γ S-ATP. AMPK complex is immuno-purified from the HEK293 cells, to which either myc-AMPK α 1 wild-type (WT) or myc-AMPK α 1 kinase-dead mutant (KD, D157A) is transfected with Flag-AMPK β 1 and HA-AMPK γ 1. AMPK complex is added into the reaction mixture containing 20 mM HEPES, pH7.4, 1 mM EGTA, 0.4 mM EDTA, 5 mM MgCl₂, 0.05 mM DTT, 0.5 μ g GST-p62, 0.2 mM AMP, and 1 mM ATP γ S. Reaction is carried out at 37°C for 30 min, and then terminated by adding 20 mM EDTA. To detect γ S-labeled p62 protein, the reaction product is alkylated with 2.5 mM PNBM for 2 h at room temperature and analyzed the products by western blotting using anti-thiophosphate antibody^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Assay ^[2]

Huh7 human hepatocytes are cultured in high glucose DMEM containing 10% FBS, 100 units/mL penicillin and 100 μ g/mL streptomycin at 37°C in a 95% air/5% CO₂ atmosphere. Hepatocytes are treated with or without Ezetimibe (10 μ M, 1 h) and incubated with palmitic acid (PA, 0.5 mM, 24 h)^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^{[1][2]}

Mice^[1]

Ten-week-old C57BL/6J male mice are used. These animals are randomly assigned to one of three groups (7-10 mice in each group): normal chow diet; MCD diet, vehicle-treated; or MCD diet, Ezetimibe -treated. The mice had free access to diet and water, with temperature maintained at 23 \pm 2°C, humidity of 60% \pm 10%, and 12-h light/dark cycles. In the MCD diet with Ezetimibe group, Ezetimibe 10 mg/kg is given once daily by oral gavage for 4 weeks. The chow and MCD diet with vehicle groups received the same volume of phosphate buffered saline orally for 4 weeks. Body weight is measured once a week over the course of the treatment period. After 4 weeks, the mice are anesthetized and killed; blood is collected via heart puncture. Tissues are harvested and either snap-frozen in liquid nitrogen and stored at -70°C or fixed in formalin and embedded in paraffin.

Rats^[2]

Male OLETF (n=11) and age-matched LETO rats (n=3) are used, and experiments are conducted in a specific pathogen-free facility with a 12 h light/dark cycle. The OLETF rat is a model that represents late-onset hyperglycemia and exhibits a chronic disease course, mild obesity and clinical onset of diabetes mellitus. Animals have unrestricted access to water and food. At 12 wk of age, rats are randomized and treated with either PBS or Ezetimibe (10 mg/kg per day) via a stomach gavage for 20 wk. At the end of the study, the rats are fasted overnight and anesthetized with intraperitoneal Zoletil/Rompun. Blood is collected from the abdominal aorta, and liver tissues are dissected, immediately frozen in liquid nitrogen, and stored at -80°C until further analysis.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- J Exp Clin Cancer Res. 2019 Sep 13;38(1):404.
- J Chem Inf Model. 2021 Jul 21.
- Front Cell Dev Biol. 18 March 2021.
- Heliyon. 2023 Oct 19.
- Virology. 2023 Jun 21.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Lee DH, et al. Ezetimibe, an NPC1L1 inhibitor, is a potent Nrf2 activator that protects mice from diet-induced nonalcoholic steatohepatitis. Free Radic Biol Med. 2016 Sep 12;99:520-532.

[2]. Chang E, et al. Ezetimibe improves hepatic steatosis in relation to autophagy in obese and diabetic rats. World J Gastroenterol. 2015 Jul 7;21(25):7754-63.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA