


Cyclophosphamide-d₈

Cat. No.:	HY-17420S1
CAS No.:	1178903-96-2
Molecular Formula:	C ₇ H ₇ D ₈ Cl ₂ N ₂ O ₂ P
Molecular Weight:	269.14
Target:	DNA Alkylator/Crosslinker; Isotope-Labeled Compounds
Pathway:	Cell Cycle/DNA Damage; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Cyclophosphamide-d ₈ is deuterium labeled Cyclophosphamide. Cyclophosphamide is a synthetic alkylating agent chemically related to the nitrogen mustards with antineoplastic activity, a immunosuppressant.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019;53(2):211-216.
- [2]. al-Jafari AA, et al. Inhibition of human acetylcholinesterase by cyclophosphamide. *Toxicology.* 1995 Jan 19;96(1):1-6.
- [3]. Harris RN, et al. Carbon tetrachloride-induced increase in the antitumor activity of cyclophosphamide in mice: a pharmacokinetic study. *Cancer Chemother Pharmacol.* 1984;12(3):167-72.
- [4]. Liu P, et al. Administration of cyclophosphamide changes the immune profile of tumor-bearing mice. *J Immunother.* 2010 Jan;33(1):53-9.
- [5]. Schwartz PS, et al. Cyclophosphamide induces caspase 9-dependent apoptosis in 9L tumor cells. *Mol Pharmacol.* 2001 Dec;60(6):1268-1279.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA