## 2'-Deoxyguanosine-<sup>13</sup>C<sub>10</sub>

Cat. No.:	HY-17563S2	
CAS No.:	1245539-86-9	HO $_{13}CH_{2}$ $H^{13}C$
Molecular Formula:	<sup>13</sup> C <sub>10</sub> H <sub>13</sub> N <sub>5</sub> O <sub>4</sub>	
Molecular Weight:	277.17	
Target:	Isotope-Labeled Compounds; Endogenous Metabolite	$\begin{array}{c c} & & & & \\ & & & \\ & & & \\ & & & \\ H^{13}C \end{array}  O_{13}CH \end{array}  N^{\prime}  NH_2$
Pathway:	Others; Metabolic Enzyme/Protease	$H^{13}C - 1^{3}CH_2$
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	ОН

<b>BIOLOGICAL ACTIV</b>	
DIOLOGICALACITY	
Description	2'-Deoxyguanosine- <sup>13</sup> C <sub>10</sub> (Deoxyguanosine- <sup>13</sup> C <sub>10</sub> ; Guanine deoxyriboside- <sup>13</sup> C <sub>10</sub> ) is <sup>13</sup> C-labeled 2'-Deoxyguanosine (HY- 17563). 2'-Deoxyguanosine (Deoxyguanosine) is deoxyguanosine.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

[2]. Deoxyguanosine, From Wikipedia

Caution: Product has not been fully validated for medical applications. For research use only.

**Product** Data Sheet



Tel: 609-228-6898 Fax: 609-2

Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA