


Crizotinib hydrochloride

Cat. No.:	HY-50878A		
CAS No.:	1415560-69-8		
Molecular Formula:	C ₂₁ H ₂₃ Cl ₃ FN ₅ O		
Molecular Weight:	486.8		
Target:	ALK; c-Met/HGFR; ROS; Autophagy		
Pathway:	Protein Tyrosine Kinase/RTK; Autophagy		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro
 H₂O : 50 mg/mL (102.71 mM; Need ultrasonic)
 DMSO : ≥ 4.9 mg/mL (10.07 mM)
 * "≥" means soluble, but saturation unknown.

	Solvent	Mass	Concentration		
			1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.0542 mL	10.2712 mL	20.5423 mL
	5 mM		0.4108 mL	2.0542 mL	4.1085 mL
	10 mM		0.2054 mL	1.0271 mL	2.0542 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description
 Crizotinib hydrochloride (PF-02341066 hydrochloride) is an orally bioavailable, selective, and ATP-competitive dual **ALK** and **c-Met** inhibitor with IC₅₀s of 20 and 8 nM, respectively. Crizotinib hydrochloride (PF-02341066 hydrochloride) inhibits tyrosine phosphorylation of NPM-ALK and tyrosine phosphorylation of c-Met with IC₅₀s of 24 and 11 nM in cell-based assays, respectively. It is also a **ROS proto-oncogene 1 (ROS1)** inhibitor. Crizotinib hydrochloride (PF-02341066 hydrochloride) has effective tumor growth inhibition^{[1][2][3]}.

IC₅₀ & Target
 IC₅₀: 20 nM (ALK), 8 nM (c-Met)^[3]

In Vitro
 PF-2341066 displays similar potency against c-Met phosphorylation in mIMCD3 mouse or MDCK canine epithelial cells with IC₅₀ of 5 nM and 20 nM, respectively. PF-2341066 shows improved or similar activity against NIH3T3 cells engineered to express c-Met ATP-binding site mutants V1092I or H1094R or the P-loop mutant M1250T with IC₅₀ of 19 nM, 2 nM and 15 nM, respectively, compared with NIH3T3 cells expressing wild-type receptor with IC₅₀ of 13 nM.

In contrast, a marked shift in potency of PF-2341066 is observed against cells engineered to express c-Met activation loop mutants Y1230C and Y1235D with IC₅₀ of 127 nM and 92 nM, respectively, compared with wild-type receptor. PF-2341066 also potently prevents the phosphorylation of c-Met in NCI-H69 and HOP92 cells, with IC₅₀ of 13 nM and 16 nM, respectively, which express the endogenous c-Met variants R988C and T1010I, respectively^[1]. PF-2341066 also potently inhibits NPM-ALK phosphorylation in Karpas299 or SU-DHL-1 ALCL cells with an IC₅₀ of 24 nM. PF-2341066 potently prevents cell proliferation, which is associated with G(1)-S-phase cell cycle arrest and induction of apoptosis in ALK-positive ALCL cells with IC₅₀ of 30 nM, but not ALK-negative lymphoma cells^[2].

In Vivo

PF-2341066 reveals the ability to cause marked regression of large established tumors (> 600 mm³) in both the 50 mg/kg/day and 75 mg/kg/day treatment cohorts, with a 60% decrease in mean tumor volume over the 43-day administration schedule in the GTL-16 model. In another study, PF-2341066 displays the ability to completely inhibit GTL-16 tumor growth for >3 months, with only 1 of 12 mice exhibiting a significant increase in tumor growth over the 3-month treatment schedule at 50 mg/kg/day. A significant dose-dependent reduction of CD31-positive endothelial cells is observed at 12.5 mg/kg/day, 25 mg/kg/day, and 50 mg/kg/day in GTL-16 tumors, indicating that inhibition of MVD shows a dose-dependent correlation to antitumor efficacy. PF-2341066 displays a significant dose-dependent reduction of human VEGFA and IL-8 plasma levels in both the GTL-16 and U87MG models. Marked inhibition of phosphorylated c-Met, Akt, Erk, PLC λ 1, and STAT5 levels is observed in GTL-16 tumors following p.o. administration of PF-2341066^[1]. Treatment of c-MET-amplified GTL-16 xenografts with 50 mg/kg PF-2341066 elicits tumor regression that is associated with a slow reduction in 18F-FDG uptake and decreases expression of the glucose transporter 1, GLUT-1^[4].

PROTOCOL

Cell Assay ^[1]

Tumor cells are seeded in 96-well plates at low density in media supplemented with 10% FBS (growth media) and transferred to serum-free media (0% FBS and 0.04% BSA) after 24 h. Appropriate controls or designated concentrations of PF-2341066 are added to each well, and cells are incubated for 24 to 72 h. Human umbilical vascular endothelial cells (HUVEC) are seeded in 96-well plates in EGM2 media for 5 to 6 h at > 20,000 cells per well and transferred to serum-free media overnight. The following day, appropriate controls or designated concentrations of PF-2341066 are added to each well, and after 1 h incubation, HGF is added to designated wells at 100 ng/mL. A 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay is done to determine the relative tumor cell or HUVEC numbers.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[1]

Athymic mice bearing xenografts (300-800 mm³) are given PF-2341066 in water by oral gavage at designated dose levels. At designated times following PF-2341066 administration, mice are humanely euthanized, and tumors are resected. Tumors are snap frozen and pulverized using a liquid nitrogen-cooled cryomortar and pestle, protein lysates are generated, and protein concentrations are determined using a BSA assay. The level of total and phosphorylated protein is determined using a capture ELISA or immunoprecipitation-immunoblotting method. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- *Cancer Discov.* 2018 Mar;8(3):354-369.
- *Nat Biomed Eng.* 2018;2:578-588.
- *Sci Transl Med.* 2018 Jul 18;10(450). pii: eaaq1093.
- *J Hematol Oncol.* 2018 Aug 29;11(1):109.

- **Cancer Res.** 2015 Nov 1;75(21):4548-59.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Zou HY, et al. An orally available small-molecule inhibitor of c-Met, PF-2341066, exhibits cytoreductive antitumor efficacy through antiproliferative and antiangiogenic mechanisms. *Cancer Res.* 2007, 67(9), 4408-4417.
- [2]. Christensen JG, et al. Cytoreductive antitumor activity of PF-2341066, a novel inhibitor of anaplastic lymphoma kinase and c-Met, in experimental models of anaplastic large-cell lymphoma. *Mol Cancer Ther.* 2007, 6(12 Pt 1), 3314-3322.
- [3]. Cui JJ, et al. Structure based drug design of crizotinib (PF-02341066), a potent and selective dual inhibitor of mesenchymal-epithelial transition factor (c-MET) kinase and anaplastic lymphoma kinase (ALK). *J Med Chem.* 2011 Sep 22;54(18):6342-63.
- [4]. Cullinane C, et al. Differential (18)F-FDG and 3'-deoxy-3'-(18)F-fluorothymidine PET responses to pharmacologic inhibition of the c-MET receptor in preclinical tumor models. *J Nucl Med.* 2011 Aug;52(8):1261-7
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA