Maleic hydrazide

Cat. No.: HY-59354

CAS No.: 123-33-1

Molecular Formula: $C_4H_4N_2O_2$

Molecular Weight: 112.09

Target: DNA/RNA Synthesis

Pathway: Cell Cycle/DNA Damage

Storage:

Powder	-20°C	3 years
	4°C	2 years
In solvent	-80°C	6 months
	-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 125 mg/mL (1115.18 mM; Need ultrasonic)

Preparing Stock Solutions

Solvent / Mass / Concentration	1 mg	5 mg	10 mg
1 mM	8.9214 mL	44.6070 mL	89.2140 mL
5 mM	1.7843 mL	8.9214 mL	17.8428 mL
10 mM	0.8921 mL	4.4607 mL	8.9214 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
   Solubility: ≥ 2.08 mg/mL (18.56 mM); Clear solution
2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
   Solubility: ≥ 2.08 mg/mL (18.56 mM); Clear solution
3. Add each solvent one by one: 10% DMSO >> 90% corn oil
   Solubility: ≥ 2.08 mg/mL (18.56 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Maleic hydrazide is extensively used as a systemic plant growth regulator and as a herbicide. Maleic hydrazide acts as an inhibitor of the synthesis of nucleic acids and proteins[1][2].

In Vitro

Maleic hydrazide is used in agriculture-in despite its known effect as a mutagenic and clastogenic agent. Maleic hydrazide had lower IC_{50} values for all cell lines compared to Ethephon. Maleic hydrazide also showed least cytotoxicity on Vero cells, followed by Hep2 and HepG2 cells[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo	Maleic hydrazide has a low acute toxicity by oral, dermal, and inhalation routes of exposure[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Venezian A, et al. The Influence of the Plant Growth Regulator Maleic Hydrazide on Egyptian Broomrape Early Developmental Stages and Its Control Efficacy in Tomato under Greenhouse and Field Conditions. Front Plant Sci. 2017;8:691. Published 2017 May 16.

[2]. Swietlińska Z, et al. Cytotoxic effects of maleic hydrazide. Mutat Res. 1978;55(1):15-30.

[3]. Yurdakok B, et al. Cytotoxic effects of etephon and maleic hydrazide in Vero, Hep2, HepG2 cells. Drug Chem Toxicol. 2014;37(4):459-465.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA