Screening Libraries

Methyl (R)-2-((tert-butoxycarbonyl)amino)-3-iodopropanoate

Cat. No.: HY-79676 CAS No.: 93267-04-0 Molecular Formula: C₉H₁₆INO₄ Molecular Weight: 329.13

Amino Acid Derivatives Target:

Pathway: Others

Storage: 4°C, protect from light

* In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)

Product Data Sheet

SOLVENT & SOLUBILITY

In Vitro

DMSO: 100 mg/mL (303.83 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	3.0383 mL	15.1916 mL	30.3831 mL
	5 mM	0.6077 mL	3.0383 mL	6.0766 mL
	10 mM	0.3038 mL	1.5192 mL	3.0383 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (7.60 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.60 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	Methyl (R)-2-((tert-butoxycarbonyl)amino)-3-iodopropanoate is an alanine derivative ^[1] .
In Vitro	Amino acids and amino acid derivatives have been commercially used as ergogenic supplements. They influence the secretion of anabolic hormones, supply of fuel during exercise, mental performance during stress related tasks and prevent exercise induced muscle damage. They are recognized to be beneficial as ergogenic dietary substances ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

1]. Luckose F, et al. Effects of an	nino acid derivatives on physica	l, mental, and physiological activ	vities. Crit Rev Food Sci Nutr. 2015;55(13):1	793-1096.
			al applications. For research use only	
	Tel: 609-228-6898 Address: 1 Dee	Fax: 609-228-5909 r Park Dr, Suite Q, Monmouth	E-mail: tech@MedChemExpress.com Junction, NJ 08852, USA	1

Page 2 of 2 www.MedChemExpress.com