


## Indalpine

Cat. No.:	HY-A0160
CAS No.:	63758-79-2
Molecular Formula:	C <sub>15</sub> H <sub>20</sub> N <sub>2</sub>
Molecular Weight:	228.33
Target:	Serotonin Transporter
Pathway:	Neuronal Signaling
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

<b>Description</b>	Indalpine (LM 5008) is a potent and selective 5-HT uptake blocker. Indalpine is potent in displacing <sup>3</sup> H-5-HT bound to brain membranes with the IC <sub>50</sub> of 36 μM <sup>[1]</sup> . Indalpine, two antidepressant agent <sup>[2]</sup> .
<b>In Vivo</b>	In keeping with its potent reuptake-blocking property, acute IV Indalpine produces a marked decrease in the firing rate of dorsal raphe 5-HT neurons (ED <sub>50</sub> 0.33 mg/kg). The firing rate of dorsal raphe 5-HT neurons is assessed following 2-, 7- and 14-day treatments with Indalpine (5 mg/day IP). After 2 days, the firing rate of 5-HT neurons is greatly reduced, after 7 days it has recovered partially and after 14 days it has returned to normal <sup>[2]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

- [1]. R Samanin, et al. Further studies on the mechanism of serotonin-dependent anorexia in rats. *Psychopharmacology (Berl)*. 1980;68(1):99-104.
- [2]. P Blier, et al. Effects of the two antidepressant drugs mianserin and Indalpine on the serotonergic system: single-cell studies in the rat. *Psychopharmacology (Berl)*. 1984;84(2):242-9.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA