


## Dipyridamole-d20

Cat. No.:	HY-B0312S		
CAS No.:	1189983-52-5		
Molecular Formula:	C <sub>24</sub> H <sub>20</sub> D <sub>20</sub> N <sub>8</sub> O <sub>4</sub>		
Molecular Weight:	524.75		
Target:	Phosphodiesterase (PDE)		
Pathway:	Metabolic Enzyme/Protease		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### BIOLOGICAL ACTIVITY

<b>Description</b>	Dipyridamole-d20 is the deuterium labeled Dipyridamole. Dipyridamole is a phosphodiesterase inhibitor that blocks uptake and metabolism of adenosine by erythrocytes and vascular endothelial cells <sup>[1][2][3]</sup> .
<b>In Vitro</b>	<p>Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

### REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother*. 2019;53(2):211-216.
- [2]. Klabunde, R.E., Dipyridamole inhibition of adenosine metabolism in human blood. *Eur J Pharmacol*, 1983. 93(1-2): p. 21-6.
- [3]. Best, L.C., et al., Mode of action of dipyridamole on human platelets. *Thromb Res*, 1979. 16(3-4): p. 367-79.
- [4]. Aktas, B., et al., Dipyridamole enhances NO/cGMP-mediated vasodilator-stimulated phosphoprotein phosphorylation and signaling in human platelets: in vitro and in vivo/ex vivo studies. *Stroke*, 2003. 34(3): p. 764-9.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA