Inhibitors

## MCE ®

## D-Glucose-d<sub>1</sub>-3

Storage:

Cat. No.: HY-B0389S7

CAS No.: 106032-60-4Molecular Formula:  $C_6H_{11}DO_6$ Molecular Weight: 181.16

Target: Endogenous Metabolite

Pathway: Metabolic Enzyme/Protease

Powder

4°C 2 years In solvent -80°C 6 months

-20°C 1 month

3 years

-20°C

## **BIOLOGICAL ACTIVITY**

| Description | D-Glucose-d-33 is the deuterium labeled D-Glucose. D-Glucose (Glucose), a monosaccharide, is an important carbohydrate in biology. D-Glucose is a carbohydrate sweetener and critical components of the general metabolism, and serve as critical signaling molecules in relation to both cellular metabolic status and biotic and abiotic stress response[1].   |
|-------------|--|
| In Vitro    | Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> .  MCE has not independently confirmed the accuracy of these methods. They are for reference only. |

## **REFERENCES**

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA