


## Ethambutol-d<sub>8</sub>

<b>Cat. No.:</b>	HY-B0535S2
<b>CAS No.:</b>	1129526-23-3
<b>Molecular Formula:</b>	C <sub>10</sub> H <sub>16</sub> D <sub>8</sub> N <sub>2</sub> O <sub>2</sub>
<b>Molecular Weight:</b>	212.36
<b>Target:</b>	Bacterial; Antibiotic; Isotope-Labeled Compounds
<b>Pathway:</b>	Anti-infection; Others
<b>Storage:</b>	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

<b>Description</b>	Ethambutol-d <sub>8</sub> is deuterium labeled Ethambutol.
<b>In Vitro</b>	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother*. 2019;53(2):211-216.
- [2]. Ethambutol. *Tuberculosis (Edinb)*, 2008. 88(2): p. 102-5.
- [3]. Kaur, D. and G.K. Khuller, In vitro, ex-vivo and in vivo activities of ethambutol and sparfloxacin alone and in combination against mycobacteria. *Int J Antimicrob Agents*, 2001. 17(1): p. 51-5.
- [4]. Rastogi, N., V. Labrousse, and K.S. Goh, In vitro activities of fourteen antimicrobial agents against drug susceptible and resistant clinical isolates of Mycobacterium tuberculosis and comparative intracellular activities against the virulent H37Rv strain in human macrophages. *Curr Microbiol*, 1996. 33(3): p. 167-75.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA