


IR 813 perchlorate

Cat. No.:	HY-D1535
CAS No.:	201024-57-9
Molecular Formula:	C ₄₀ H ₄₀ Cl ₂ N ₂ O ₄
Molecular Weight:	683.66
Target:	Others
Pathway:	Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	IR 813 perchlorate is a near-infrared (NIR) fluorescent dye (Ex=815 nm, Em=840 nm) and can be used for visualizing regional lymph nodes in mice ^[1] .
In Vitro	IR 813 (17.3 μM, 2 h) induces a 31.4% hemolysis in red blood cells ^[1] . IR 813 (5.9 μM, 24 h) induces 50% cell death in MRC-5 cells ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	IR 813 perchlorate displays a maximum fluorescence intensity at 4 h postinjection together with a rapid extravasation, when being used for visualizing regional lymph nodes in mice ^[1] . Guidelines (Following is our recommended protocol. This protocol only provides a guideline, and should be modified according to your specific needs) ^[1] . 1. Animal: Ten-week-old female Balb/cOlaAnN mice are kept in a 12 h light/dark cycle to reduce tissue autofluorescence in the NIR region, and had access to food and water ad libitum. 2. Dose: A single dose 5.1 nmol of IR 813 dye (20 μL of 0.173 mg/mL, dissolved in a PEG-400/ethanol/water=3:2:5, v/v/v solution) is subcutaneous injected in the right anterior paw of mice. 3. Imaging: Using the Fluobeam700 NIR imaging system to perform in vivo optical imaging (739 nm excitation light (3.5 mW), 750 nm long-pass emission cutoff filter). Fluorescence intensity in the axillary lymph node (ALN) is recorded for 1 week (5 min, 1 h, 4 h, 24 h, 7 days). 4. Data analysis: Semiquantitative data is obtained from the fluorescence images using ImageJ 1.44 software. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Marion Hell, et al. Surface chemistry architecture of silica nanoparticles determine the efficiency of in vivo fluorescence lymph node mapping. ACS Nano. 2013 Oct 22;7(10):8645-57.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA