


## CM-H2DCFDA

<b>Cat. No.:</b>	HY-D1713
<b>CAS No.:</b>	850013-49-9
<b>Molecular Formula:</b>	C <sub>27</sub> H <sub>19</sub> Cl <sub>3</sub> O <sub>8</sub>
<b>Molecular Weight:</b>	577.79
<b>Target:</b>	Reactive Oxygen Species
<b>Pathway:</b>	Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κB
<b>Storage:</b>	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



### BIOLOGICAL ACTIVITY

<b>Description</b>	CM-H2DCFDA is a derivative of <a href="#">H2DCFDA</a> (HY-D0940). CM-H2DCFDA can be used to determine cellular oxidant levels (Ex/Em: 495/530 nm). CM-H2DCFDA is light-sensitive <sup>[1]</sup> .
<b>In Vitro</b>	Guidelines <sup>[2]</sup> (Following is our recommended protocol. This protocol only provides a guideline, and should be modified according to your specific needs). ROS Measurement: 1. Incubate the cells according to your normal protocol. 2. Treat cells with 10 μM CM-H2DCFDA for 30 min at 37°C in darkness. 3. Wash the excess probe. 4. Analyze sample on a flow cytometer, fluorescence microscopy, or fluorescence microplate reader. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### CUSTOMER VALIDATION

- Front Public Health. 2023 Jul 13;11:1222762.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

### REFERENCES

[1]. Oparka M, et al. Quantifying ROS levels using CM-H2DCFDA and HyPer. *Methods*. 2016 Oct 15;109:3-11.

[2]. Kolarova H, et al. Production of reactive oxygen species after photodynamic therapy by porphyrin sensitizers. *Gen Physiol Biophys*. 2008 Jun;27(2):101-5.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA