


## 18β-Glycyrrhetic acid

<b>Cat. No.:</b>	HY-N0180		
<b>CAS No.:</b>	471-53-4		
<b>Molecular Formula:</b>	C <sub>30</sub> H <sub>46</sub> O <sub>4</sub>		
<b>Molecular Weight:</b>	470.68		
<b>Target:</b>	Endogenous Metabolite		
<b>Pathway:</b>	Metabolic Enzyme/Protease		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	1 year
		-20°C	6 months



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 125 mg/mL (265.57 mM; Need ultrasonic)					
		Solvent Concentration	Mass	1 mg	5 mg	10 mg
	<b>Preparing Stock Solutions</b>	1 mM		2.1246 mL	10.6229 mL	21.2459 mL
		5 mM		0.4249 mL	2.1246 mL	4.2492 mL
10 mM			0.2125 mL	1.0623 mL	2.1246 mL	
Please refer to the solubility information to select the appropriate solvent.						
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.17 mg/mL (4.61 mM); Clear solution  2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.17 mg/mL (4.61 mM); Clear solution					

### BIOLOGICAL ACTIVITY

<b>Description</b>	18β-Glycyrrhetic acid is the major bioactive component of Glycyrrhiza uralensis and possesses anti-ulcerative, anti-inflammatory and antiproliferative properties.
<b>In Vitro</b>	18β-Glycyrrhetic acid is the major bioactive component of Glycyrrhizae Radix and possesses anti-ulcerative, anti-inflammatory and antiproliferative properties. MTS assay demonstrates that 24 h treatment of 18β-Glycyrrhetic acid suppresses cell proliferation in both cell lines in a dose-dependent manner. 18β-Glycyrrhetic acid at 160 μM significantly decreases the percentage of viable cells to around 40.5±10.5% in A549 and 38.3±4.6% in NCI-H460 (p<0.01 respectively). When the cells are treated with 320 μM 18β-Glycyrrhetic acid, a greater inhibitory effects on cell proliferation is shown, as the percentage of viable cells is below 30% compare with untreated controls (p<0.001). Treatment with 18β-Glycyrrhetic acid at 160 μM and 320 μM decreases the levels of full-length PARP and increases the levels of cleaved-PARP <sup>[1]</sup> .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

Rats in 18 $\beta$ -Glycyrrhetic acid+Triptolide (TP) group which receive low-dose 18 $\beta$ -Glycyrrhetic acid (50 mg/kg) have significant reductions in the three serum parameters when compare with TP rats. Rats in 18 $\beta$ -Glycyrrhetic acid+TP group which receive the high-dose 18 $\beta$ -Glycyrrhetic acid (100 mg/kg) have slightly lowered the levels of three liver enzymes, the reductions do not reach statistical significance compare with TP group. Contrastingly, preadministration of low-dose 18 $\beta$ -Glycyrrhetic acid protects animals from TP-induced hepatic lesions. On the contrary, low-dose 18 $\beta$ -Glycyrrhetic acid (50 mg/kg) markedly suppresses the release of the four cytokines above<sup>[3]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

#### Cell Assay <sup>[2]</sup>

Primary microglia cultures are used in this study. For treatment assay, microglia are incubated with complete DMEM and stimulated with or without 100 ng/mL IFN- $\gamma$  in the presence or absence of 18 $\beta$ -Glycyrrhetic acid (25  $\mu$ M and 50  $\mu$ M) at 37°C in a humidified incubator with 5% CO<sub>2</sub>. For cell migration assay, the isolated primary microglia that seeded in complete DMEM medium are stimulated with or without IFN- $\gamma$  (100 ng/mL), and treated with different doses of 18 $\beta$ -Glycyrrhetic acid, 24 h later, the microglia culture supernatants are collected and added to the lower chambers of Transwell inserts<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Animal Administration <sup>[3]</sup>

Healthy Wistar rats (male, 200 $\pm$ 20 g) are used and divided into five groups with 10 individuals for each group randomly. Animals in normal control (NC) group receive distilled water for 6 days and 0.5% CMC-Na for the last 3 days. Rats in Triptolide model group (TP), 18 $\beta$ -Glycyrrhetic acid low-dose group (GAL+TP), and 18 $\beta$ -Glycyrrhetic acid high-dose group (GAH+TP) receive distilled water, 18 $\beta$ -Glycyrrhetic acid (50 mg/kg, p.o., dissolved in distilled water), or 18 $\beta$ -Glycyrrhetic acid (100 mg/kg, p.o., dissolved in distilled water) for consecutive 6 days, respectively, and liver injury is induced by TP (2.4 mg/kg, p.o., suspended in 0.5% CMC-Na) for the last 3 days. Animals in the above three groups receive TP 6 hours after distilled water or 18 $\beta$ -Glycyrrhetic acid treatment on the last 3 days<sup>[3]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Adv Sci (Weinh). 2024 Jan 6:e2305260.
- Chemosphere. 2023 Feb 24;138249.
- Cell Prolif. 2023 May 4;e13494.
- Environ Toxicol. 2022 Aug 3.
- World J Gastroenterol. 2023 Jun 21; 29(23): 3622-3644.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Huang RY, et al. 18 $\beta$ -Glycyrrhetic acid suppresses cell proliferation through inhibiting thromboxane synthase in non-small cell lung cancer. PLoS One. 2014 Apr 2;9(4):e93690.

[2]. Zhou J, et al. 18 $\beta$ -glycyrrhetic acid suppresses experimental autoimmune encephalomyelitis through inhibition of microglia activation and promotion of remyelination. Sci Rep. 2015 Sep 2;5:13713.

[3]. Yang G, et al. Protective Effect of 18 $\beta$ -Glycyrrhetic Acid against Triptolide-Induced Hepatotoxicity in Rats. Evid Based Complement Alternat Med. 2017;2017:3470320.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA