


Benzoic acid lithium

Cat. No.:	HY-N0216A
CAS No.:	553-54-8
Molecular Formula:	C ₇ H ₅ LiO ₂
Molecular Weight:	128.05
Target:	Bacterial; Fungal; Endogenous Metabolite
Pathway:	Anti-infection; Metabolic Enzyme/Protease
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

H₂O : 100 mg/mL (780.94 mM; Need ultrasonic)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	7.8094 mL	39.0472 mL	78.0945 mL
	5 mM	1.5619 mL	7.8094 mL	15.6189 mL
	10 mM	0.7809 mL	3.9047 mL	7.8094 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Benzoic acid lithium is an aromatic alcohol found in many plants and is a common additive in food, beverages, cosmetics and other products. Benzoic acid lithium inhibits bacteria and fungi and acts as a preservative^[1].

In Vitro

Benzoic acid (1 mM) lithium stimulates OAT2-catalyzed efflux of glutamic acid and orotic acid from OAT2h-expressing 293 cells^[2].

Benzoic acid (0.5 and 2.0 mg/disc) lithium shows antibacterial activity against *S. aureus*, *S. epidermidis*, *P. aeruginosa*, *B. cereu*)^[4].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Combinations of benzoic acid (2,000 mg/kg in diet, 1-14 days) lithium and Thymol (HY-N6810) (100 mg/kg in diet, 1-14 days) improves growth performance, promotes nutrient digestion and absorption, and reduces diarrhea in weaned pigs^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA