## **Product** Data Sheet

# L-Methionine-<sup>15</sup>N

Cat. No.:HY-N0326SCAS No.:82572-25-6Molecular Formula: $C_5H_{11}^{115}NO_2S$ Molecular Weight:150.2

Target:Endogenous MetabolitePathway:Metabolic Enzyme/Protease

Storage: Powder -20°C 3 years 4°C 2 years

In solvent  $-80^{\circ}$ C 6 months  $-20^{\circ}$ C 1 month

$$S \longrightarrow OH$$

$$15NH_2$$

#### **SOLVENT & SOLUBILITY**

In Vitro

 $H_2O: \ge 50 \text{ mg/mL} (332.89 \text{ mM})$ 

\* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	6.6578 mL	33.2889 mL	66.5779 mL
	5 mM	1.3316 mL	6.6578 mL	13.3156 mL
	10 mM	0.6658 mL	3.3289 mL	6.6578 mL

Please refer to the solubility information to select the appropriate solvent.

### **BIOLOGICAL ACTIVITY**

**Description**L-Methionine-<sup>15</sup>N is the <sup>15</sup>N-labeled L-Methionine. L-Methionine is the L-isomer of Methionine, an essential amino acid for human development. Methionine acts as a hepatoprotectant.

In Vitro Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### **REFERENCES**

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$ 

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com