## L-Glutamine-<sup>13</sup>C<sub>5</sub>

Cat. No.:	HY-N0390S	1		
CAS No.:	184161-19-	1		
Molecular Formula:	<sup>13</sup> C <sub>5</sub> H <sub>10</sub> N <sub>2</sub> O <sub>3</sub>			O L O
Molecular Weight:	151.11			$H_2N^{-13}C_{-$
Target:	mGluR; Fer	roptosis;	Endogenous Metabolite	$H_2$ $13C$
Pathway:	GPCR/G Pro	otein; Ne	uronal Signaling; Apoptosis; Metabolic Enzyme/Protease	NH <sub>2</sub>
Storage:	Powder	-20°C	3 years	
		4°C	2 years	
	In solvent	-80°C	6 months	
		-20°C	1 month	

## SOLVENT & SOLUBILITY

	Concentration	1 mg	5 mg	10 mg			
Preparing Stock Solutions	1 mM	6.6177 mL	33.0885 mL	66.1770 mL			
	5 mM	1.3235 mL	6.6177 mL	13.2354 mL			
	10 mM	0.6618 mL	3.3088 mL	6.6177 mL			
Please refer to the solubility information to select the appropriate solvent.							
	Preparing Stock Solutions Please refer to the sol	Preparing       1 mM         Stock Solutions       5 mM         10 mM       10 mM         Please refer to the solubility information to select the approximation to select the	Preparing Stock Solutions       1 mM       6.6177 mL         5 mM       1.3235 mL         10 mM       0.6618 mL         Please refer to the solubility information to select the appropriate solvent.	Preparing Stock Solutions       1 mM       6.6177 mL       33.0885 mL         5 mM       1.3235 mL       6.6177 mL         10 mM       0.6618 mL       3.3088 mL         Please refer to the solubility information to select the appropriate solvent.			

DIOLOGICAL ACTIV					
Description	L-Glutamine- <sup>13</sup> C <sub>5</sub> is the <sup>13</sup> C-labeled L-Glutamine. L-Glutamine (L-Glutamic acid 5-amide) is a non-essential amino acid present abundantly throughout the body and involved in many metabolic processes. L-Glutamine provides a source of carbons for oxidation in some cells[1][2].				
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.				

## REFERENCES



[1]. Mary Corless, et al. Glutamine Regulates Expression of Key Transcription Factor, Signal Transduction, Metabolic Gene, and Protein Expression in a Clonal Pancreatic Beta-Cell Line. J Endocrinol. 2006 Sep;190(3):719-27.

[2]. Newsholme P, et al. Glutamine and glutamate as vital metabolites. Braz J Med Biol Res. 2003 Feb;36(2):153-63. Epub 2003 Jan 29.

[3]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

## Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA