


Phillygenin

Cat. No.:	HY-N0483
CAS No.:	487-39-8
Molecular Formula:	C ₂₁ H ₂₄ O ₆
Molecular Weight:	372.41
Target:	Reactive Oxygen Species
Pathway:	Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κB
Storage:	4°C, protect from light, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light, stored under nitrogen)



SOLVENT & SOLUBILITY

In Vitro

DMSO : 50 mg/mL (134.26 mM; Need ultrasonic)
H₂O : < 0.1 mg/mL (insoluble)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	2.6852 mL	13.4261 mL	26.8521 mL
	5 mM	0.5370 mL	2.6852 mL	5.3704 mL
	10 mM	0.2685 mL	1.3426 mL	2.6852 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 3.25 mg/mL (8.73 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 3.25 mg/mL (8.73 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 3.25 mg/mL (8.73 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Phillygenin (Phillygenol) is an active ingredient from Forsythia with many medicinal properties, such as antioxidant, reducing blood lipid, inhibition of low density lipoprotein oxidation.

In Vitro

- Phillygenin shows a greater inhibition on mouse B16 melanoma cells potential than vincristine^[1].
- phillygenin had notable scavenging activity against DPPH, ABTS radicals, as well as potent reducing power in FRAP assay. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

The reference for rat is 5.6 mg/ml (i.v).

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Biosci BioTech Bioch. 2021 Jan 9.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Ye LH et al. Determination of phillygenin in rat plasma by high-performance liquid chromatography and its application to pharmacokinetic studies. Eur J Drug Metab Pharmacokinet, 2013 Sep, 38(3):201-7.

[2]. Song W et al. Interaction between phillygenin and human serum albumin based on spectroscopic and molecular docking. Spectrochim Acta A Mol Biomol Spectrosc, 2012 Jan, 85(1):120-6.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA