## **Product** Data Sheet

## L-Leucine-<sup>13</sup>C<sub>6</sub>, <sup>15</sup>N

Cat. No.: HY-N0486S8

CAS No.: 202406-52-8

Molecular Formula: 13C<sub>6</sub>H<sub>13</sub>15NO<sub>2</sub>

Molecular Weight: 138.12

Target: mTOR; Endogenous Metabolite


Pathway: PI3K/Akt/mTOR; Metabolic Enzyme/Protease

**Storage:** Powder -20°C 3 years

4°C 2 years -80°C 6 month

In solvent -80°C 6 months

-20°C 1 month



## **BIOLOGICAL ACTIVITY**

Description	L-Leucine- $^{13}$ C <sub>6</sub> , $^{15}$ N is the $^{13}$ C- and $^{15}$ N-labeled L-Leucine. L-Leucine is an essential branched-chain amino acid (BCAA), which activates the mTOR signaling pathway[1].
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> .  MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## **REFERENCES**

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Inhibitors