


trans-Isoferulic acid

Cat. No.:	HY-N0761A
CAS No.:	25522-33-2
Molecular Formula:	C ₁₀ H ₁₀ O ₄
Molecular Weight:	194.18
Target:	NO Synthase; Prostaglandin Receptor
Pathway:	Immunology/Inflammation; GPCR/G Protein
Storage:	4°C, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)



BIOLOGICAL ACTIVITY

Description	trans-Isoferulic acid (trans-3-Hydroxy-4-methoxycinnamic acid) is an aromatic acid isolated from the roots of <i>Clematis florida</i> var. <i>plena</i> . trans-Isoferulic acid exhibits anti-inflammatory activity ^[1] , trans-isoferulic acid suppresses NO and PGE2 production through the induction of Nrf2-dependent heme oxygenase-1 (HO-1) ^[2] .
IC ₅₀ & Target	Human Endogenous Metabolite

REFERENCES

[1]. Kai-Hui Sun, et al. A new indole-type alkaloid from the roots of *Clematis florida* var. *plena*. *Nat Prod Res.* 2019 Oct;33(20):2925-2931.

[2]. Matharage Gayani Dilshara, et al. Downregulation of NO and PGE2 in LPS-stimulated BV2 microglial cells by trans-isoferulic acid via suppression of PI3K/Akt-dependent NF-κB and activation of Nrf2-mediated HO-1. *Int Immunopharmacol.* 2014 Jan;18(1):203-11.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA