


Catalpol

Cat. No.:	HY-N0820		
CAS No.:	2415-24-9		
Molecular Formula:	C ₁₅ H ₂₂ O ₁₀		
Molecular Weight:	362.33		
Target:	HBV		
Pathway:	Anti-infection		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 30 mg/mL (82.80 mM)
 * "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	2.7599 mL	13.7996 mL	27.5992 mL
	5 mM	0.5520 mL	2.7599 mL	5.5198 mL
	10 mM	0.2760 mL	1.3800 mL	2.7599 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: ≥ 2.08 mg/mL (5.74 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
 Solubility: ≥ 2.08 mg/mL (5.74 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
 Solubility: ≥ 2.08 mg/mL (5.74 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Catalpol (Catalpinoside), an iridoid glycoside found in *Rehmannia glutinosa*. Catalpol has neuroprotective, hypoglycemic, anti-inflammatory, anti-cancer, anti-spasmodic, anti-oxidant effects and anti-HBV effects^{[1][2][3]}.

In Vitro

Catalpol (0.1 μg/mL; for 3 days) can induce neuronal differentiation in PC12 cells through activation of the intracellular signal transduction pathway, and promote neurite length^[2].
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Catalpol (25-100 mg/kg; intraperitoneal injection; once) treatment clearly reduces blood urea nitrogen, serum creatinine levels and the expression of KIM-1 in renal ischemia/reperfusion-injury (IRI) mice. Catalpol markedly reduces the expression of PI3K, Akt and eNOS levels, and suppresses the TNF- α , IL-1 β , IL-6 and IL-10 activities^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	C57BL/6 mice treated with renal ischemia/reperfusion surgery ^[1]
Dosage:	25 mg/kg, 50 mg/kg and 100 mg/kg
Administration:	Intraperitoneal injection; once
Result:	Clearly reduced blood urea nitrogen, serum creatinine levels and the expression of KIM-1 in renal IRI mice.

CUSTOMER VALIDATION

- Pharmacol Res. 2020 May;155:104751.
- Acta Pharmacol Sin. 2021 Nov 18.
- Phytomedicine. 2020 Nov;78:153300.
- Microvasc Res. 2021 Dec 14;104302.
- SSRN. 2021 Oct.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Jili Zhu, et al. Catalpol protects mice against renal ischemia/reperfusion injury via suppressing PI3K/Akt-eNOS signaling and inflammation. *Int J Clin Exp Med*. 2015 Feb 15;8(2):2038-44.

[2]. M Yamazaki, et al. Neuritogenic effect of natural iridoid compounds on PC12h cells and its possible relation to signaling protein kinases. *Biol Pharm Bull*. 1996 Jun;19(6):791-5.

[3]. R Mehrotra, et al. In vitro studies on the effect of certain natural products against hepatitis B virus. *Indian J Med Res*. 1990 Apr;92:133-8.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA