Product Data Sheet

Thymidine-¹³C₁₀

Cat. No.: HY-N1150S11

Molecular Formula: ${}^{13}C_{10}H_{14}N_2O_5$

Molecular Weight: 252.16

Target: Isotope-Labeled Compounds; DNA/RNA Synthesis; Endogenous Metabolite;

Orthopoxvirus

Pathway: Others; Cell Cycle/DNA Damage; Metabolic Enzyme/Protease; Anti-infection

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.

$$H_3^{13}C_{13}C_{\overset{13}{\circ}C}$$
 NH
$$H^{13}C_{\overset{13}{\circ}C}$$
 NH
$$H^{13}C_{\overset{13}{\circ}C}$$

$$H^{13}C_{\overset{13}{\circ}C}$$

$$H^{13}C_{\overset{13}{\circ}C}$$

$$H^{13}C_{\overset{13}{\circ}C}$$

$$H^{13}C_{\overset{13}{\circ}C}$$

$$H^{13}C_{\overset{13}{\circ}C}$$

BIOLOGICAL ACTIVITY

Description	Thymidine- 13 C ₁₀ (DThyd- 13 C ₁₀ ; NSC 21548- 13 C ₁₀) is 13 C-labeled Thymidine (HY-N1150). Thymidine, a specific precursor of deoxyribonucleic acid, is used as a cell synchronizing agent. Thymidine is a DNA synthesis inhibitor that can arrest cell at G1/S boundary, prior to DNA replication.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.
- [2]. Chen G, et al. Cell Synchronization by Double Thymidine Block. Bio Protoc. 2018 Sep 5;8(17).
- [3]. FIRKET H, et al. Autoradiographic visualization of synthesis of deoxyribonucleic acid in tissue culture with tritium-labelled thymidine. Nature. 1958 Jan 24;181(4604):274-5. FIRKET H, et al. Autoradiographic visualization of synthesis of deoxyribonucleic acid in tissue culture with tritium-labelled thymidine. Nature. 1958 Jan 24;181(4604):274-5.
- [4]. Izeradjene K, et al. Inhibition of thymidine synthesis by folate analogues induces a Fas-Fas ligand-independentdeletion of superantigen-reactive peripheral T cells. Int Immunol. 2001 Jan;13(1):85-93.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA