## Linocinnamarin

MedChemExpress

| Cat. No.:<br>CAS No.:<br>Molecular Formula:<br>Molecular Weight:<br>Target:<br>Pathway:<br>Storage: | HY-N11887<br>554-87-0<br>C <sub>16</sub> H <sub>20</sub> O <sub>8</sub><br>340.33<br>Reactive Oxygen Species<br>Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κB<br>Please store the product under the recommended conditions in the Certificate of<br>Analysis. |  |
|---|---|--|
|---|---|--|

| DI | $\mathbf{n}$ | $\mathbf{n}$ | 210 | A I    | ΛС | · T I \ | VIT۱ | / |
|----|--------------|--------------|-----|--------|----|---------|------|---|
| Ы  | UL           | $\mathbf{U}$ |     | ÷1 - 7 | нu |         | V    |   |
|    |              |              |     |        |    |         |      |   |

Description Linocinnamarin is a nature product that could be isolated from Fragaria ananassa Duch. Linocinnamarin has antiinflammatory activity. Linocinnamarin inhibits antigen-stimulated elevation of intracellular free Ca<sup>2+</sup> concentration and reactive oxygen species (ROS)<sup>[1]</sup>.

## REFERENCES

[1]. Ninomiya M, et, al. Phenolic constituents isolated from Fragaria ananassa Duch. inhibit antigen-stimulated degranulation through direct inhibition of spleen tyrosine kinase activation. Bioorg Med Chem. 2010 Aug 15;18(16):5932-7.

Caution: Product has not been fully validated for medical applications. For research use only.

Fax: 609-228-5909 Tel: 609-228-6898 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

**Product** Data Sheet