## Myristic acid-d<sub>2</sub>

Cat. No.: CAS No.: Molecular Formula: Molecular Weight: Target: Pathway: Storage:	HY-N2041S1 30719-21-2 C <sub>14</sub> H <sub>26</sub> D <sub>2</sub> O <sub>2</sub> 230.38 Endogenous Metabolite; Isotope-Labeled Compounds Metabolic Enzyme/Protease; Others Please store the product under the recommended conditions in the Certificate of	он р р он
	Analysis.	

BIOLOGICAL ACTIV		
BIOLOGICALACIWIT		
Description	Myristic acid-d <sub>2</sub> is the deuterium labeled Myristic acid. Myristic acid is a saturated 14-carbon fatty acid occurring in most animal and vegetable fats, particularly butterfat and coconut, palm, and nutmeg oils.	
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

## REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Inhibitors

•

**Screening Libraries** 

•

Proteins