


2 α ,19 α -Dihydroxy-3-oxo-urs-12-en-28-oic acid

Cat. No.:	HY-N4155
CAS No.:	176983-21-4
Molecular Formula:	C ₃₀ H ₄₆ O ₅
Molecular Weight:	486.68
Target:	HIV Protease; HSV; EBV
Pathway:	Anti-infection; Metabolic Enzyme/Protease
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	2 α ,19 α -Dihydroxy-3-oxo-urs-12-en-28-oic acid, a natural ursane-type triterpene, is a potent inhibitor of HIV protease (HIV Protease). 2 α ,19 α -Dihydroxy-3-oxo-urs-12-en-28-oic acid is also an inhibitor of the activation of Epstein-Barr virus early antigen (EBV-EA). 2 α ,19 α -Dihydroxy-3-oxo-urs-12-en-28-oic acid displays an inhibitory activity against nitric oxide production in Lipopolysaccharide (Lipopolysaccharides)-activated RAW 264.7 cells ^{[1][2]} .								
In Vitro	2 α ,19 α -Dihydroxy-3-oxo-urs-12-en-28-oic acid (0.01-0.1 μ M) displays moderate inhibitory activities against nitric oxide production in Lipopolysaccharide-activated macrophage cell lines, RAW 264.7 cells ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.								
In Vivo	2 α ,19 α -Dihydroxy-3-oxo-urs-12-en-28-oic acid (0.0025%; drinking water; for 20 weeks) shows an inhibitory effect on the activation of EBV-EA induced by TPA and causes a significant delay of two-stage carcinogenesis on mouse skin ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.								
	<table border="1"> <tr> <td>Animal Model:</td> <td>Female ICR mice (6 weeks old)^[2]</td> </tr> <tr> <td>Dosage:</td> <td>0.0025%</td> </tr> <tr> <td>Administration:</td> <td>Drinking water; for 20 weeks</td> </tr> <tr> <td>Result:</td> <td>Showed an inhibitory effect on the activation of EBV-EA induced by 12-O-tetradecanoylphorbol-13-acetate (TPA).</td> </tr> </table>	Animal Model:	Female ICR mice (6 weeks old) ^[2]	Dosage:	0.0025%	Administration:	Drinking water; for 20 weeks	Result:	Showed an inhibitory effect on the activation of EBV-EA induced by 12-O-tetradecanoylphorbol-13-acetate (TPA).
Animal Model:	Female ICR mice (6 weeks old) ^[2]								
Dosage:	0.0025%								
Administration:	Drinking water; for 20 weeks								
Result:	Showed an inhibitory effect on the activation of EBV-EA induced by 12-O-tetradecanoylphorbol-13-acetate (TPA).								

REFERENCES

- [1]. Xiao-Peng Wu, et al. A new ursane-type triterpene, cymosic acid from *Rosa cymosa*. *J Asian Nat Prod Res.* 2014;16(4):422-5.
- [2]. Shoko Taniguchi, et al. Production of bioactive triterpenes by *Eriobotrya japonica* calli. *Phytochemistry.* 2002 Feb;59(3):315-23.
- [3]. Xiao-Peng Wu, et al. A new ursane-type triterpene, cymosic acid from *Rosa cymosa*. *J Asian Nat Prod Res.* 2014;16(4):422-5.
- [4]. Shoko Taniguchi, et al. Production of bioactive triterpenes by *Eriobotrya japonica* calli. *Phytochemistry.* 2002 Feb;59(3):315-23.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA