


## Mogroside IIe

Cat. No.:	HY-N6814
CAS No.:	88901-38-6
Molecular Formula:	C <sub>42</sub> H <sub>72</sub> O <sub>14</sub>
Molecular Weight:	801.01
Target:	Others
Pathway:	Others
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



### SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (124.84 mM; Need ultrasonic)						
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg	
				1 mM	1.2484 mL	6.2421 mL	12.4842 mL
				5 mM	0.2497 mL	1.2484 mL	2.4968 mL
				10 mM	0.1248 mL	0.6242 mL	1.2484 mL
Please refer to the solubility information to select the appropriate solvent.							
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (3.12 mM); Clear solution						
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (3.12 mM); Clear solution						
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (3.12 mM); Clear solution						

### BIOLOGICAL ACTIVITY

Description	Mogroside IIe is a nonsugar sweetener. Mogrosides are sweeter than sucrose. Mogrosides exhibit antioxidant, antidiabetic and anticancer activities <sup>[1]</sup> .
-------------	---

### REFERENCES

[1]. Itkin M, et al. The biosynthetic pathway of the nonsugar, high-intensity sweetener mogroside V from *Siraitia grosvenorii*. Proc Natl Acad Sci U S A. 2016 Nov 22;113(47):E7619-E7628.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA