

Grape seed extract

Cat. No.:	HY-N7072
CAS No.:	84929-27-1
Target:	Apoptosis
Pathway:	Apoptosis
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)

Grape seed extract

SOLVENT & SOLUBILITY

In Vitro	DMSO : 6 mg/mL (Need ultrasonic)
In Vivo	<ol style="list-style-type: none"> Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 0.6 mg/mL (Infinity mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 0.6 mg/mL (Infinity mM); Clear solution

BIOLOGICAL ACTIVITY

Description	Grape seed extract is a natural product, with anti-inflammatory and anti-proliferative effects. Grape seed extract shows inhibitory activity on the fat-metabolizing enzymes pancreatic lipase and lipoprotein lipase. Grape seed extract induces apoptotic in human colorectal cancer cells ^{[1][2][3]} .										
In Vitro	<p>Grape seed extract (25-100 µg/mL; 12-48 hours) inhibits growth and induces death in human colorectal cancer cells^[3]. Grape seed extract (25-100 µg/mL; 12-24 hours) modulates the protein levels of cell cycle regulatory molecules in HT29 and LoVo cells^[3]. Grape seed extract (25-100 µg/mL; 24 hours) causes apoptotic cell death of human colorectal cancer cells^[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Cell Proliferation Assay^[3]</p> <table> <tr> <td>Cell Line:</td> <td>HT29 cells, LoVo cells</td> </tr> <tr> <td>Concentration:</td> <td>25 µg/mL, 50 µg/mL, 100 µg/mL</td> </tr> <tr> <td>Incubation Time:</td> <td>12 hours, 24 hours, 48 hours</td> </tr> <tr> <td>Result:</td> <td>Inhibited cell growth in a dose- and a time-dependent manner.</td> </tr> </table> <p>Cell Cycle Analysis^[3]</p> <table> <tr> <td>Cell Line:</td> <td>HT29 cells, LoVo cells</td> </tr> </table>	Cell Line:	HT29 cells, LoVo cells	Concentration:	25 µg/mL, 50 µg/mL, 100 µg/mL	Incubation Time:	12 hours, 24 hours, 48 hours	Result:	Inhibited cell growth in a dose- and a time-dependent manner.	Cell Line:	HT29 cells, LoVo cells
Cell Line:	HT29 cells, LoVo cells										
Concentration:	25 µg/mL, 50 µg/mL, 100 µg/mL										
Incubation Time:	12 hours, 24 hours, 48 hours										
Result:	Inhibited cell growth in a dose- and a time-dependent manner.										
Cell Line:	HT29 cells, LoVo cells										

	<table border="1"> <tr> <td>Concentration:</td> <td>25 µg/mL, 50 µg/mL, 100 µg/mL</td> </tr> <tr> <td>Incubation Time:</td> <td>12 hours, 24 hours, 48 hours</td> </tr> <tr> <td>Result:</td> <td>Induced cell cycle arrest (a marked increase in Cip1/p21 protein level and a decrease in G 1 phase associated cyclins and cyclin-dependent kinases).</td> </tr> </table>	Concentration:	25 µg/mL, 50 µg/mL, 100 µg/mL	Incubation Time:	12 hours, 24 hours, 48 hours	Result:	Induced cell cycle arrest (a marked increase in Cip1/p21 protein level and a decrease in G 1 phase associated cyclins and cyclin-dependent kinases).		
Concentration:	25 µg/mL, 50 µg/mL, 100 µg/mL								
Incubation Time:	12 hours, 24 hours, 48 hours								
Result:	Induced cell cycle arrest (a marked increase in Cip1/p21 protein level and a decrease in G 1 phase associated cyclins and cyclin-dependent kinases).								
	<p>Western Blot Analysis^[3]</p> <table border="1"> <tr> <td>Cell Line:</td> <td>HT29 cells, LoVo cells</td> </tr> <tr> <td>Concentration:</td> <td>25 µg/mL, 50 µg/mL, 100 µg/mL</td> </tr> <tr> <td>Incubation Time:</td> <td>12 hours, 24 hours</td> </tr> <tr> <td>Result:</td> <td>Modulates the protein levels of CDKIs, CDKs, and cyclins.</td> </tr> </table>	Cell Line:	HT29 cells, LoVo cells	Concentration:	25 µg/mL, 50 µg/mL, 100 µg/mL	Incubation Time:	12 hours, 24 hours	Result:	Modulates the protein levels of CDKIs, CDKs, and cyclins.
Cell Line:	HT29 cells, LoVo cells								
Concentration:	25 µg/mL, 50 µg/mL, 100 µg/mL								
Incubation Time:	12 hours, 24 hours								
Result:	Modulates the protein levels of CDKIs, CDKs, and cyclins.								
	<p>Apoptosis Analysis^[3]</p> <table border="1"> <tr> <td>Cell Line:</td> <td>HT29 cells, LoVo cells</td> </tr> <tr> <td>Concentration:</td> <td>25 µg/mL, 50 µg/mL, 100 µg/mL</td> </tr> <tr> <td>Incubation Time:</td> <td>24 hours</td> </tr> <tr> <td>Result:</td> <td>Showed a significant dose-dependent increase in apoptotic cell population in HT29 cells at 50 µg/mL and 100 µg/mL; all three Concentrations caused a dose-dependent increase in apoptotic cells.</td> </tr> </table>	Cell Line:	HT29 cells, LoVo cells	Concentration:	25 µg/mL, 50 µg/mL, 100 µg/mL	Incubation Time:	24 hours	Result:	Showed a significant dose-dependent increase in apoptotic cell population in HT29 cells at 50 µg/mL and 100 µg/mL; all three Concentrations caused a dose-dependent increase in apoptotic cells.
Cell Line:	HT29 cells, LoVo cells								
Concentration:	25 µg/mL, 50 µg/mL, 100 µg/mL								
Incubation Time:	24 hours								
Result:	Showed a significant dose-dependent increase in apoptotic cell population in HT29 cells at 50 µg/mL and 100 µg/mL; all three Concentrations caused a dose-dependent increase in apoptotic cells.								
In Vivo	<p>Grape seed extract (200 mg/kg; i.g.; 5 days/wk; for 8 weeks) inhibits HT29 colon carcinoma tumor xenograft growth in athymic nude mice^[3].</p> <p>Grape seed extract inhibits cell proliferation but increases apoptotic cell death in tumors^[3].</p> <p>Grape seed extract enhances Cip1/p21 protein levels and poly(ADP-ribose) polymerase cleavage in tumors^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1"> <tr> <td>Animal Model:</td> <td>Athymic male nu/nu mice, with human colon carcinoma HT29 tumor xenograft^[3]</td> </tr> <tr> <td>Dosage:</td> <td>200 mg/kg</td> </tr> <tr> <td>Administration:</td> <td>Oral gavage, 5 days/wk, for 8 weeks</td> </tr> <tr> <td>Result:</td> <td>Showed time-dependent inhibition of tumor growth.</td> </tr> </table>	Animal Model:	Athymic male nu/nu mice, with human colon carcinoma HT29 tumor xenograft ^[3]	Dosage:	200 mg/kg	Administration:	Oral gavage, 5 days/wk, for 8 weeks	Result:	Showed time-dependent inhibition of tumor growth.
Animal Model:	Athymic male nu/nu mice, with human colon carcinoma HT29 tumor xenograft ^[3]								
Dosage:	200 mg/kg								
Administration:	Oral gavage, 5 days/wk, for 8 weeks								
Result:	Showed time-dependent inhibition of tumor growth.								

CUSTOMER VALIDATION

- Int J Mol Sci. 2023 Apr 6, 24(7), 6826.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. The effects of grape seed extract (Vitis vinifera) supplement on inflammatory markers, neuropeptide Y, anthropometric measures, and appetite in obese or overweight

individuals: A randomized clinical trial.

[2]. Inhibitory effects of grape seed extract on lipases. Nutrition. 2003 Oct;19(10):876-9.

[3]. Manjinder Kaur, et al. Grape seed extract inhibits in vitro and in vivo growth of human colorectal carcinoma cells. Clin Cancer Res. 2006 Oct 15;12(20 Pt 1):6194-202.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA